

Abstracts of the Papers Presented at the 24th Biennial Conference of the AABE

The 24th Biennial Conference of the AABE was held at University of the Philippines, Diliman, Quezon City, Philippines, from 5th to 9th December, 2012. The theme of the Conference was “The Century of Biology: Towards Transformative Education.” Nearly 150 people from ten countries including outside of the Asian region attended (Figure 1).



Figure 1 Attendants assembling in front of the conference venue.

There were five plenary lectures, four country reports, 32 oral presentations and 34 poster presentations. In addition, three workshops were presented.

The best presentation awards were given to the following papers:

(Oral presentation)

Diong, C. H. & Oh, S. S. Importance of Parasite Diversity in the Understanding of Global Biodiversity.

Obusan, M. C. M. Where Dolphins Are.

(Poster presentation)

Yamanoi, T., Iwasaki, W. M., Takemura, M. & Sakura, O. Origami Bird: A Teaching Material Linking Mutation, Natural Selection and Speciation.

Chiga, S. & Ohshika, K. Development of Teaching Material Using Owl Pellet of the Zoo.

The Book of the Abstracts of Papers presented at this conference was published and provided to the attendants of the conference. The following abstracts are reprinted from the book.

<Plenary Lectures>

The Asian Biodiversity Crisis: The Other Crisis in Asia

Perry S. Ong

University of the Philippines, Philippines

Asia is one of the most biodiversity rich regions in the world. Six of the 17 megadiversity countries are found in the region: the Philippines, China, India, Malaysia, Indonesia and Papua New Guinea, while nine out of 34 biodiversity hotspots are from Asia.

These six countries comprise nearly 40% of global population, which exerts tremendous pressure to transform these biodiversity resources into material wealth that will lift their people out of poverty. The impact of one of the drivers of biodiversity loss is being felt across the region.

To illustrate this twin distinction of pride and grief, of the 633 species of primates all over the world, about

54% are under various categories of threat. Of course, the top 25 endangered primate species had been identified, of which nine species are from Asia: Vietnam = five species, Indonesia = three species, and China = one species. This means Asia is home to 36% of the top 25 endangered primates in the world.

Thus, the loss of biodiversity and the need to conserve this valuable resource is a common thread across the region. One of the reasons for the loss is the low level of awareness about the rich biodiversity the region possesses. The role of education, in particular the Asian Association for Biology Education (AABE), cannot be over emphasized. While the Internet is a powerful tool to assist in educating the general public, we should ensure that science-based information is communicated. The key message for all to disseminate is that “A healthy, living and thriving biodiversity is the foundation of a sustainable future for Asia and the rest of the world.”

Perry S. Ong, Ph.D., Institute of Biology, College of Science, UP-Diliman, Quezon City, Philippines

From *Gessellschaftlich* to *Gemeinschaft*: What Does It Takes to Teach with and for Meaning in Biology?

Allan B De Guzman

University of Santo Tomas, Philippines

The changing landscape of today's society brought about by forces of consumerism, globalization, environmental degradation, poverty, economic inequalities, illiteracy, and poor health delivery services, among others poses a challenge to basic and higher education sectors to adopt a transformative pedagogy through which learners are reunited with their environment in the name of learning excellence and in the spirit of genuine service. This emerging pedagogy puts premium on the reciprocal relationship between student centrality and community instrumentality in developing an authentic way of knowing (cognitive), an experiential way of doing (conative) and a more reflective way of responding (affective).

Anchored on Ballard's triad of the change process, which consists of (1) *awareness* of what is happening and what is required; (2) *agency* or ability to find a response that seems personally meaningful; and (3) association with other people in groups and networks, this paper is an attempt to couple service learning or education in action and sustainability as a means to pedagogize the existing teaching-learning paradigm in today's education system. As a multi-tiered pedagogy that can be implemented at any level of education, pre-school through graduate school, service learning (SL) or community engaged teaching (CET) recognizes and appreciates the value of the community as responsive and effective partners in students' learning and maintains the centrality of students as active players and contributors to sustainable development. In the final analysis, pedagogy of sustainability (PS) becomes a critical variable in defining not only the proactiveness of today's schools, colleges and universities but more importantly as an effective compass in asserting education's leadership presence and extraordinariness toward the creation of a common future through an integrated teaching, research and extension

service framework anchored on service learning.

Allan B. De Guzman, Ph.D. abdeguzman@mnl.ust.edu.ph

Education Transformation and Intel STEM

Yvonne Garcia Flores

Intel Philippines

Intel advocates the systemic approach to transforming education, providing interventions that acknowledge and address issues of the whole education environment: Policy, Curriculum and Assessment, Professional Development, Research and Evaluation, and Information and Communications Technology. This particular presentation will showcase Intel's latest education resources specific to Science, Technology, Engineering and Math (Intel STEM), such teacher professional development, online tools and platforms.

Does Postmodern Ethics Have Anything to Say to the Sciences as Presently Constituted?

Fr. Luis S. David, S.J.

Ateneo de Manila University, Philippines

The social sciences emerged in the 19th century, on Michael Foucault's account, in the confluence of two orders of knowledge, one, traditional, going all the way back to Plato and Aristotle, that decidedly was intuitive, inductive, experiential, etc., and the other, springing from the principal philosophies of modernity, including those of Descartes, Rousseau, and Hobbes, which was conjectural, speculative, suggestive of what Hannah Arendt calls, "a science of prophecy." The latter practice, which underservedly wraps itself around with the mantle of scientific legitimacy borrowed from the former, arguably has led to totalitarianisms and genocidal events. We face the challenge, therefore, of reinforcing the claims of the former, and insisting that the latter, when unavoidable, sooner rather than later receive "traction" through its correlation with inductive work. This requires that we "work on ourselves," that in our communities of scientific practice we remain grounded in what is observable, intersubjective, etc., that our

work be understood as always “in process,” “in progress,” etc., subject to the A to Z of what is “political” and therefore communitarian.

An Interdisciplinary Approach to Biology Education

Lourdes J. Cruz

University of the Philippines, and President, National Research Council of the Philippines, Philippines

Biology, the study of living organisms interfaces with other disciplines of science. Understanding the structure, function, growth, origin, evolution and distribution of organisms requires application of the principles of chemistry, physics, mathematics, geology and other fields. An interdisciplinary approach is necessary for the students to fully understand the principles of biology and to learn how living organisms relate to the physical and social world around us. Students must be well equipped to face the problems that confront us today such as food security, sustainability, and climate change because these problems will become more serious as we pass on the baton to the younger generation. To illustrate how an interdisciplinary approach can be used in biology education, aquaponics will be discussed as an example of the interplay of organisms in an ecological system and the importance of chemical and physical factors in sustainably maintaining the system. The backyard aquaponics system has a great potential as a means for poor families to regularly have food on their table. If expanded, the system can be used as an alternative means of livelihood.

Lourdes J. Cruz, Ph.D., Marine Science Institute, UP-Diliman, Quezon City, Philippines

<Country Reports>

Biology Education in Australia

Anne M Wallis

Deakin University, Australia

An Australian national curriculum covering “Foundation schooling”(preparatory year) through to Year 10 has recently been introduced into Australian schools. Previously, each Australian states or territory had responsibility for school curricula, funding and overall

educational policy. States and territories still have responsibility for the final two years of school years – Years 11 and 12 which usually cover ages 16 to 18.

Science forms one of the 13 key *learning areas* in what is now known as the Australian Curriculum. Each learning area is based on seven generic capabilities relating to Literacy, Numeracy, Information and Communication Technology Capability, Critical and Creative Thinking, Ethical Behaviour, Personal and Social Capability, and Intercultural Understanding, and three cross-curriculum properties (Aboriginal and Torres Strait Islander histories and cultures, Sustainability and Asia, and Australia’s engagement with Asia). The 13 year Science curriculum has four strands –one of which is Biological Science which covers evolution, ecology, organism structure and function as well as consideration of life’s major groups.

In my own State of Victoria, Year 11 and 12 are covered by a range of qualifications, the most popular of which is the Victorian Certificate of Education (VCE). Students typically take English (compulsory) and four other subjects in their final year. Biology is still the most popular of the core sciences (and in 2011 was the 7th most popular subject), although its popularity is stable as students seek more applied and focussed subjects, such as Health and Human Development which last year had over 1500 more enrolments and Psychology had over 3000 more enrolments than Biology. The VCE Biology Curriculum has been updated and modified program of study will commence in 2013. The four units to be taken (covering two years of Biology study) are

Unit 1: Unity and Diversity

Unit 2: Organisms and Their Environment

Unit 3: Signatures of Life

Unit 4: Continuity and Change

The unit titles and most content is the same as that reported by me in the 2008 Country Report, but assessment has been changed to better reflect the expected student outcomes.

In universities, Biology is still a popular and often core component in science degrees. However, in many applied degrees, science educators are becoming increasingly concerned at the fragmentation of content and learning in Biology, so that while graduates may have detailed expertise in specific areas, they might

totally lack fundamental knowledge in others that provide the building blocks for the application of Biology to areas such as health or environmental science. One of the greatest challenges for educators today will be the transition of Cloud Learning.

Dr. Anne M. Wallis, School of Life and Sciences, Deakin University, Warrnambool, Vic. 3290, Australia

Biological Learning Contents for High School Students in Textbooks According to 2009 National Science Curriculum of Korea

Kyoungho Kim¹⁾, Kew-Cheol Shim²⁾

¹Gongju National University of Education, ²Kongju National University, South Korea

New national science curriculum of Korea was revised in 2009. On the basis of that, new science textbooks were developed for high school students. High school science textbooks are Physics I, II, Chemistry I, II, Biological Science I, II Earth Science I, II and Science, which is so called 'Fusing Science' or 'Convergence Science.' Objectives of them consist of four domains regarding as cognitive, inquiry, affective, and STS ones. High school students have to select and learn some subjects for more than 15 credits. Science of the 2009 revised curriculum has two unites related to biological sciences such as 'Evolution of Life' and 'Human Health and Technology.' Biological Science I has 'Understanding of Biological Sciences,' 'Cell and Continuity of Lives,' 'Homeostasis and Health,' and 'Human in the Nature.' Biological Science II has 'Cell and Metabolism,' 'Gene and Bioengineering,' and 'Evolution of Organisms.'

Professor Kyoungho Kim, kkh@gjue.ac.kr; Professor Kew-Cheol Shim, skcshim@kongju.ac.kr

New Biology Education in Japan: Characteristics of the New Course of Study and New Program for Biology Teachers in Japan

Ohshika Kiyoyuki

Aichi University of Education, Japan

The New Science Curriculum which based on the revised Course of Study prescribed by the ministry of Education, Culture, Sports, Science and Technology in

2008 and 2009 had started in all school levels from elementary to senior high school in Japan. The New Science Curriculum focuses on the systematical formation and enrichment of scientific contents, inquiry-based activities with emphasis on experimentations and observation, and real life experience.

However, there are some problems in carrying out biology education under the New Science Curriculum. For example, time is considered as a limiting factor for teachers to teach biology contents which it require. In the New Science Curriculum, time element is so short to teach the subject matter. Most of the young teachers have not enough confidences to teach biological content knowledge, conduct classroom experiments and outdoor experiences because of their background in the former biology curriculum during their baccalaureate courses and they had little exposure to biological content knowledge and pedagogical skills. Furthermore, the New Biology Curriculum adds some new contents at each school level. For example, the topics on invertebrates, non-vascular plants and evolution were transferred to junior high school biology; DNA and genetic engineering were added. Therefore, it is suggested to develop and formulate many programs for teachers and students in biology education. All teachers were required to take workshops or lectures in 30 hours to update their own teacher certification every ten years.

To support science classes, there are special lectures offered through the Science Partnership Program and Super Science High School Program that support schools and universities or specialists in science. This undertaking will lead to the development of the newest and creative biology education in Japan through these policies and programs.

Dr. Kiyoyuki Ohshika, Aich University of Education, Hirosawa 1, Igaya-cho, Kariya, Aich 448-8542, Japan

<Oral presentations>

Where Dolphins Are

Marie Christine Merca Obusan

University of the Philippines, Philippines

Space use models were developed for spinner dolphins (*Stenellalongirostris*) in Tañon Strait, Philippines.

Home ranges were generated based on kernel density, minimum convex polygon, Jennrich-Turner primary axis, site fidelity test, and spider analysis using the Animal Movement Program (version 2) as an extension of Arcview. Analyses of developed models validated the need to consider seasonal patterns in designing management strategies for the conservation and protection of the cetacean species in site. The spatial and temporal information provided can serve as basis for associating the dynamics between dolphin survival and human activities such as local fishing and tourism.

Prof. Marie Christine Merca Obusan, Institute of Environmental Science and Meteorology, College of Science, UP-Diliman, Quezon City; Philippine Science High School - Bicol Region Campus, Goa, Camarines Sur, Philippines

Multipla-loci cpDNA Cladistic Frameworks Reveal Polyphyly of *Antirhea* Juss (Guettardeae-Rubiaceae) and a Circumscription of a New Philippine Endemic

Jayson Chavez

Far Eastern University, Philippines

In the last comprehensive synopsis of the paleotropical *Antirhea* Juss., the genus which distribution is highly concentrated in the Philippines is divided into three subgenera: *Antirhea*, *Guettardella* and *Mesocarpa*; a classification proposed primarily based on classical taxonomy and up until now, is a subject of disputes among systematics. For the present study, a molecular phylogeny of the genus was constructed for the first time with the following objectives: (1) test the monophyly of *Antirhea* and its relationships to related genera, (2) evaluate the subgenera of *Antirhea* proposed by Chaw and Darwin, and (3) determine the identity of an unusual *Antirhea* species collected during the recent botanical expedition of the Thomasian Angiosperm Phylogeny and Barcoding Group (TAPBG). Sequences of two plasmid markers (*rps16* and *trnL-F*) were analyzed separately as well as combined using parsimony (MP) and Bayesian (BI) approaches. Our results highly support the polyphyly of *Antirhea* as currently circumscribed, but further support the monophyly of *Antirhea* subgenus *Guettardella*. Such inferences reveal disparities with the phonetic classifica-

tion of the genus. Extensive investigations on *Antirhea*, which involve a wide-ranging survey (e.g., denser taxon sampling, thorough morphological examinations, and cladistic analyses of DNA markers with high evolutionary rate) are currently ongoing to delimit its taxonomic boundaries. In addition, based on herbarium and field study, the atypical *Antirhea* species is treated as a new endemic species: *A. acuminata* Di-matatac, Chavez & Alejandro, which is proposed here.

Some Easy and Effective Microtechniques for Morpho-anatomical Characterization of Plants: An Essential Tool in Transformative Learning

Vivian S. Tolentino, Jan Lorie M. Robil

Ateneo de Manila University, Philippines

Morphology and anatomy are indispensable aspects that are employed to study and extensively characterize our flora and fauna. It is an essential tool for documentation of plant species, especially those which are newly discovered or understudied, as it gives relevant information about their physiology, ecology and life history. Thus, knowledge in microtechniques is indispensable for both researchers and educators of plant biology. However, tools and special equipment for microtechniques are not always readily available. Expensive microtomes that are used for cutting thin tissue sections are not always available especially in schools in rural areas. In this paper, easy and effective microtechniques are discussed, using two Philippine indigenous plant species, *Medinilla magnifica* (chandelier plant) and *Strongylodon macrobotrys* (jade vine) as models. This paper aims to give information on how to process plant tissue sections without the use of expensive cutting equipment, so as to practically aid researchers and educators in their study of indigenous plants. Free-hand sectioning method was used for the vegetative structures of both specimens. Prior to cutting, the specimens have undergone fixation in formalin-acetic acid-alcohol (FAA). Portions of leaves and stems were sectioned manually using razor blades to produce very thin sections of tissues. These sections were then subjected to series of dehydration using ethyl alcohol (30%, 50%, 70%, 90%, and 100% EtOH).

The sections were then stained and counterstained using Safranin and Fast Green, respectively. The tissues were washed with water and were destained in 70% EtOH and were subjected to series of cleaning using xylene (30%, 50%, 70%, 90%, and 100% xylene). Finally, the cleared sections were mounted in glass slides. The whole procedure which took only about an hour produced mounted tissue sections which can be used for characterization of ground and vascular tissue structures of vegetative organs of the specimens. This easy and rapid tissue processing is an essential tool for teaching biology and research, hence making the laboratory activities more enjoyable and fun.

Prof. Vivian S. Tolentino, Department of Biology, School of Science and Engineering, Ateneo de Manila University, Loyola Hights, Quezon City 1108, Philippines

The Importance of Understanding Relationships among Biological Species Using a Phylogenetic Tree

Grecebio Jonathan Alejandro
University of Santo Tomas, Philippines

A phylogenetic tree (also known as evolutionary tree or a cladogram) is a branching diagram depicting the relationships and affiliations among biological species. The taxa included to build a “tree” are joined together based on either morphological features or genetic characteristics or combination of the two. The resulting tree reflects that the taxa have descended from a common ancestor. The arrangement of biological species in a phylogenetic tree is easier for the students to understand and interpret relationships. In this lecture, an introduction to phylogenetics will be presented. The concept of groupings approaches terminologies and methodologies in phylogenetics will be outlined. The usefulness of molecular data in phylogenetics will be discussed using examples from the studies done by the Thomasian Angiosperm Phylogeny and Barcoding Group (TAPBG).

Dr. Grecebio Jonathan Alejandro, College of Science and Research Centre for the Natural and Applied Sciences, Thomas Aquinas Research Complex, University of Santo Tomas, Philippines

Compliance Status to CHED Policies and Standards on Teacher Education and Institutional Performance of Teacher Education Institutions in Region IX

Mario Rojas Obra, Jr., Carmelita M. Felisilda
Western Mindanao State University, Philippines

This study purported to create an accurate picture of the teacher education discipline in the light of CHED Policies and Standards on Teacher Education and bring to the fore the institutional performance of Teacher Education Institutions (TEIs) in Region IX. Specifically, the study endeavored to answer the following research questions: (1) What is the compliance status of TEIs in Region IX to CHED Policies and Standards, in terms of program administration-dean, faculty, physical facilities, laboratory facilities, research, admission and retention, library holdings, voluntary accreditation, curriculum, and instructional standards; (2) Is there a significant difference in the compliance status of TEIs in Region IX to CHED Policies and Standards when data are grouped according to type of TEIs (public or private); (3) What is the institutional performance of TEIs in Region IX in terms of passing percentage in Licensure Examinations for Teachers (LET) for elementary and secondary levels and frequency in landing among the Top Ten schools in LET; (4) Is there a significant difference in the institutional performance to TEIs in Region IX when data are grouped according to type of TEIs (public or private).

When data are grouped according to type of TEIs, it was hypothesized that (1) there is a significant difference in the compliance status of TEIs in Region IX to CHED Policies and Standards in terms of the ten given areas of academic operations and (2) there is a significant difference in the institutional performance of TEIs in Region IX in terms of the passing percentage in LET.

The study employed descriptive research design and used qualitative and quantitative approach in answering the set of research questions. Such process was facilitated with the use of a researcher-made interview schedule/guide which helped generate data inputs from the responses/interview proceedings made to 71 purposively selected middle level administrators, faculty and staff coming from the ten participating TEIs in

Region IX. Proceedings from actual interview, analyses of documentary evidences and data-output from statistical tools used revealed the following findings: TEIs in Region IX showed full compliance on six academic operations of CHED Policies and Standards: administration-dean (2.7), faculty (2.4), physical facilities (2.76), admission and retention (2.4), curriculum (3.0) and instructional standards (2.52), but manifested partial compliance on four other academic operations, to wit: laboratory facilities (1/95), research (1.95), library holdings (1.92) and voluntary accreditation (1.8). Inferential findings of the study revealed higher group mean of public TEIs (2.55) relative to private TEIs (2.26). Moreover, t-test analysis showed a probability value of 0.72 greater than alpha level 0.05, that implies no significant difference in the compliance status of TEIs in Region IX to CHED Policies and Standards. Latter finding showed that the type of school does not influence the TEIs adherence to CHED Policies and Standards. Both public and private TEIs have similar compliance status to CHED Policies and Standards. Moreover, the type of school does not influence the institutional performance of TEIs in terms of passing percentage in LET for elementary and secondary levels. In addition, the performance standing of TEIs in Region IX commensurate national recognition as evidenced by its frequency in landing among the Top Ten performing schools in LET for elementary and secondary levels during the fiscal year 2008 and 2010. It is, therefore, recommended for CHED Regional Office IX to conduct periodic standard monitoring with a show of report to feedback TEIs for self evaluation and improvement; and for school's top management to foresee enhancement of personal and professional competencies of the faculty and improve the inadequacies of school facilities, equipments, and library holdings, while middle level management must do periodic review of the curriculum to see subject offerings appropriateness; improve the efficiency level of the faculty along areas of research, extension and production; and for other research enthusiasts to integrated TEI graduates employability as one of the independent variables to get clearer and better picture of its institutional performance.

Prof. Mario Rojas Obra, Jr., College of Education, Western Mindanao State University, Zamboanga City, Philippines

Secondary Teachers and Pre-service Teachers' Conceptual Understanding of Photosynthesis: A Cross Regional Study

**Jocelyn D. Partosa¹, Michael A. Clores²,
Maria Aurora A. Conde², Maricar S. Prudente³,
Lydia T. Goingo², Arnulfo R. Reganit²**

¹Ateneo de Zamboanga University, ²Ateneo de Naga University, ³De La Salle University, Philippines

This study aimed to identify misconceptions that secondary in-service teachers and pre-service teachers have of photosynthesis and compare their knowledge structure of photosynthesis. The misconceptions of in-service and pre-service secondary teachers from conveniently selected high schools in Regions V and IX were identified using Treagust and Haslam's Two-Tiered Diagnostic test on Photosynthesis and Cellular Respiration (1987). Interviews with randomly selected secondary and pre-service teachers were done to determine the knowledge structure of the respondents. Generally, for both secondary in-service and pre-service teachers, the misconceptions revolve on any one or a combination of the following: failure to differentiate between photosynthesis and respiration, failure to give a functional understanding of the chemical basis of biological processes and for some, failure to correctly identify the site of biological processes. The respondents have shallow and flawed understanding of both photosynthesis and cellular respiration. The inability to explain and show how the foregoing concepts relate is more pronounced among the pre-service teachers. The foregoing misconceptions point to faulty, fragmentary and incomplete knowledge structures across all respondents. Thus, the conceptual understanding is at the knowledge level only for most pre-service teachers.

Prof. Jocelyn D. Partosa, Natural Science Department, Ateneo de Zamboanga University, Philippines

Incorporation of Mother Tongue in Biology Subjects: Preference among Philippine Science High School Central Visayas Campus Students

Sherry Ramayla

Philippine Science High School - Central Visayas Campus, Philippines

This paper is based on the mandate of the Department of Education to use the mother tongue in primary schools as medium of instruction. The goal of this study is to determine language preferences of the students of Philippine Science High School – Central Visayas Campus, Argao, Cebu, in their Biology classes. A survey was conducted June 28, 2012, using structured questionnaires. The sample consisted of 22 second year students, 22 third year students, and 26 fourth year students. The respondents were randomly selected. Findings showed that majority of the respondent preferred bilingual medium of instruction (English and the mother tongue in Cebu, Cebuano). Students find the combination of English and Cebuano as conversational or it encourages classroom interaction between teacher and student or student and student and it allows the students to grasp more of the meaning of the concept of Biology if the two languages are used inside the classroom.

Prof. Sherry Ramayla, Philippine Science High School - Central Visayas Campus, Cebu, Philippines

Secondary Teachers' Biological Conceptions of Natural Selection: A Cross Regional Study

**Michael A. Clores¹, Jocelyn D. Partosa²,
Maria Aurora A. Conde¹, Maricar S. Prudente³,
Lydia T. Goingo¹, Arnulfo R. Reganit¹**

¹Ateneo de Naga University, ²Ateneo de Zamboanga University, ³De La Salle University, Philippines

This mixed method research determined the conceptual understanding of pre-service and in-service secondary biology teachers in Bicol and Zamboanga about 'Natural Selection' by generating: level of understanding, knowledge structures, alternative conceptions and misconceptions. A total of 113 respondents took the Conceptual Inventory of Natural Selection in the first part of the study. In the second part, interviews of 37 were analyzed using NVivo9 and content analysis. The level of understanding of teachers about the Natural Selection is generally *low*; they found most of the test items to be *difficult*, and they expressed numerous alternative conceptions and misconceptions. Seven dominant alternative conceptions were identified and are affirmed by four nodes, namely, "Need," "Use and

disuses," "Progress," and "Purpose." Based on the results, recommendation on instruction, curriculum and research were forwarded.

Dr. Michael A. Clores, Natural Science Department, Ateneo de Naga University; Prof. Jocelyn D. Partosa, Natural Science Department, Ateneo de Zamboanga University, Philippines

The Silkworm as a Teaching Material

Koichi Morimoto

Nara University of Education, Japan

The silkworm is well-known and has already been used as a teaching material for a long time widely in Japan. As the silkworm is very useful for teaching of several contents, it is expected that it will be good teaching material in Asian countries other than Japan.

In primary school biology, the silkworm is suitable for the observation of the life cycle of insect. It has egg, larva, pupa and adult stages. Students can observe how it feeds in the mulberry leaves and how it makes a cocoon, and they can also observe the mating behavior of it. The students can discover many things in magnified image.

In lower secondary school biology, the silkworm is useful for the observation of structure of insect in the Arthropod. Students can observe the spiracle dorsal vessel, the nail of legs and the simple eyes. The mating behavior is also very attractive for the students.

In upper secondary school biology, students can understand the role of the silk gland which produces the silk. As the silkworm vomits the digestive juice when it receives an electric shock, the students can confirm the function of the digestive juice by using it. One of the phenotypes of the silkworm is the markings of its body. The students can confirm Mendel's Law by using the phenotype.

Active Contents-based Learning in Science Education

Kazutoshi Tamura

Niigata Municipal Bandai Senior High School, Japan

New contents-based learning plan is implemented in two years. The plan is the contents-based learning incorporated with inquiry-based learning. We call the plan of learning "Active contents-based learning."

We suggest that the method of the inquiry-based learning is attractive and effective when it is used as the part of the contents-based learning. Inquiry-based learning and contents-based learning are complementary each other. Therefore, a problem of these methods is that how to unite the methods of inquiry-based learning and of the contents-based learning. In Japan, education of biology is cling to think about merely description of organism's phenomenon, not enough to think about principles of life and to gain the idea of organisms systems, thus mainly done the contents-based learning. Therefore, Active contents-based learning is valid in Japan.

From this year, 2012, senior secondary school curriculum guidelines are changed in Japan. New contents are increased in textbooks. Also, inquiry-based learning is recommended by the guidelines, because the OECD test scores (PISA) in all areas have been gradually decreased since 2003. However, inquiry-based learning method is not enough increased. One reason is that inquiry-based learning becomes the solitary new one subject. The other reason is that high school teachers do not understand enough the inquiry-based learning. In Japan, high school education is relatively poor to gain the creativity, to try new action in science and also many areas in culture. High school education is needed to change such the matter in Japan. However, it is difficult to alter the cultural situation. Fortunately, our nation has also somewhat abundant history of science research. Biology education and science education will be changed gradually like to our plan, "Active inquiry-based learning."

Prof. Kazutoshi Tamura, Niigata Municipal Bandai Senior High School, Niigata, Japan

An Innovative Approach in Teaching Total RNA Isolation in Genetics, Cell and Molecular Biology Laboratory

Vivian Panes

Ateneo de Manila University, Philippines

The method of isolating DNA from various types of organisms, in particular using a formulated and modified DNA isolation protocol instead of a commercial kit, is oftentimes introduced in laboratory classes in Genetics, as well as in Cell and Molecular Biology in

the Philippines. The method of isolating RNA, however, is rarely introduced in these classes except in the thesis of collegiate level biology students in some universities in the country. The preference for introducing DNA isolation rather than RNA is obvious, in that the DNA, being double stranded is more stable than RNA and hence the outcome for DNA extraction is oftentimes guaranteed relative to RNA, specifically with the use of a laboratory concocted isolation protocol.

The use of commercial kits for RNA isolation is certainly very expensive. Consequently, the high cost of commercial kits for RNA isolation is a limitation to many universities in the Philippines. Hence, the aim of this paper is to present an innovative, less expensive method for total RNA isolation. The method basically focuses on total RNA isolation of indigenous Philippine plants with valuable commercial and medicinal attributes, such as *Artemisia vulgaris* (damong Maria) and *Zingiber officinale* (ginger). A modified procedure for isolating total RNA from a mushroom, *Ganoderma lucidum* is also introduced in this paper. The formulation of this modified RNA isolation method enabled the students to become more creative and become less dependent on kits. Although kits have advantages, such kits can serve as technical crutches that dull the students' ability to interpret experiments and understand where they have gone wrong, and may lose their ability to troubleshoot effectively.

Prof. Vivian Panes, Department of Biology, School of Science and Engineering, Ateneo de Manila University, Loyola Heights, Quezon City, Philippines

Transmission Electron Microscopy of In Vitro Biofilms Formed by *Candida albicans* and *Escherichia coli*

Frederick Masangkay

PAMET, PASMETH, PSM, MIMLS, Philippines

Bacteria and *Candida* may adhere to surfaces or grow in matrix-enclosed biofilms. This study investigated biofilm architecture of *Candida albicans* and *Escherichia coli* sessile cells in pure and co-cultures *in vitro*.

Significance: How biofilms are formed is key to understanding effects in the human host when pathogenic

organisms: *Candida albicans* associated with yeast infections and *Escherichia coli* the most common cause of gastro-enteritis and urinary tract infection establishes a community. Interaction between these organisms would demonstrate phenotypic characteristics which will serve as basis in discovering appropriate strategies and interventions particular to these microbial communities.

Methods: Clinical isolates of bacterial and fungal cells were grown in thioglycollate broth. Sessile cells (biofilm phenotype) were harvested by decanting the bulk fluid, swabbing the slides of the test tubes and planting in BAP, MAC and CAP. Sessile cells were then grown on copper grids for 72 hours and negatively stained with 2% phosphotungstic acid for Transmission Electron Microscopy examination. Sessile cells were also grown in Eppendorf tubes with thioglycollate for 72 hours, fixed with glutaraldehyde and processed for positive staining. Ultra-thin sections were stained with 7% methanolic uranyl acetate solution and lead citrate.

Results and Discussion: *C. albicans* sessile cells in pure and co-culture with *E. coli* in thioglycollate did not produce hyphae or pseudohyphae; it maintained the blastospore (yeast cell) morphology. Biofilm architecture of towers predominated in *C. albicans* pure culture and demonstrated the formation of “blobs.” Networking architecture was observed in pure colonies of *E. coli*. Negative staining was superior in demonstrating exopolysaccharide structures because of absence of dehydration process. However, positive staining was superior in demonstrating cell walls. Sessile cells of *E. coli* and *C. albicans* in *in vitro* co-cultures in thioglycollate broth resulted in down-regulated biofilm formation demonstrated by decreased cellularity of biofilms and degradation of yeast cells.

Retrial to Select Biological Terms for Secondary School Science Education in Japan

Nobuyasu Katayama

Tokyo Institute of Biology Education, Japan

As a result of the latest reform of the national curricu-

lum standard, the Course of Study (CS), by the Ministry of Education, Culture, Sports, Science and Technology, Japan (MEXT), the contents of biology-related subjects have significantly changed. In particular, the CS for upper secondary schools has been very much modernized. In addition, the relaxation of the textbook screening process by MEXT has resulted in a wide variety of synonymous biological terms appearing in biology textbooks which are published by some private publishers. This is particularly remarkable in the upper secondary school level. Upper secondary school students, therefore, must remember all of these terms if they will take university entrance examinations. It must be a burden on these students. The Society of Biological Sciences Education Japan (SBSEJ) has organized a working group (the SBSEJ Examination Committee for Biological Terms) for selecting biological terms appearing in biology textbooks for upper secondary schools to reduce the burden on the students and for revising the Glossary of Biological Terms which was published in 1998. In the presentation, we will report our activities to date. We also show some data such as the number of biological terms appearing in the glossary section of Basic Biology - Biology for All - textbooks which were published last March and the redundancy of some of these terms among these textbooks.

Dr. Nobuyasu Katayama, nobukei@oregano.ocn.ne.jp

Scientific Researches among Articles of Scientifically Gifted Students for Biology Education

**Ji-Hyon Kil¹⁾, Kew-Cheol Shim²⁾, Kyoungho Kim³⁾,
Jung-Min Kim⁴⁾**

^{1)National Institute of Environmental Research, ^{2)Kongju National University, ^{3)Gongju National University of Education, ^{4)Environmental Ecological Education Institute, South Korea}}}}

This study is to analyze articles of scientifically gifted students, which were published in Korean Journal of Young Scientist. They were written by scientifically gifted students who were high school students. The papers were reviewed by scientists and most of them were research papers resulted from scientific research-based mentoring programs. The scientific mentoring

programs were performed by high school students with being helped by university professors or institute scientists. Their research themes were related to mathematics, physics, chemistry, biological sciences and earth sciences. Articles of them related to biological sciences showed in experimental methods and tools including DNA and protein works. The results implied that various laboratory experiences motivated and facilitated them to understand biological properties, and writing an article should be useful to facilitate eceintific ability of scientifically gifted students.

Prof. Kyoungho Kim, Department of Science Education, Gongju National University of Education, Gongju, Chungnam, South Korea

Utilization of Learner-centered Principles in Grade 4 and 5 Math and Science Instructional Activities in a Jesuit School in Metro Manila

Mel Greg Conception
Xavier School, Philippines

In any Jesuit school, *cura personalis* means having personal care for the students. The learners serve as the focus of the academic program and institution. Researchers claim that students increase their motivation sand achievement if teachers provide them with utmost care and positive learning environment. This research aims to study what and how academic subject teachers in Math and Science utilize to make their institutions and teaching practices learner-centered. The methodology used in this research was a qualitative descriptive research. Participants were randomly selected using fishbowl sampling technique from a total of 576 grade 4 and 5 students. The students' perception was measured by a simple student questionnaire on how their teachers' use of learner- centered principles affects their motivation and academic performance. Selection of teacher participants were grade school math and science teachers and were categorized as novice, junior and senior teachers. Grades 4 and 5 math and science teacher participants were asked to answer the questionnaire. Likewise, they were observed in their classes by their respective department head. Documentary analysis was utilized to gather data from the classroom observations conducted by the

department heads. Focus was on the leamer- centered instructional activities provided by the math and science teachers in this study. Correction was established with the students' motivation and academic performance and the teachers' use of learner-centered principles in math and science subjects specifically on instructional activities.

Prof. Mel Greg Conception, Middle School Science Department, Xavier School, Greenhills West, San Juan City; Department of Biological Sciences, Philippine Normal University, Manila, Philippines

Development and Application of Concept Based Objective Tests (CBOTs) for High School Students' Understanding of the School Biology Syllabus

Narendra D. Deshmukh
Homi Bhabha Centre for Science Education, TIFR, India

Identification and characterizing of misconception generally involve considerable effort. Different tools are used by different researchers to study the effectiveness of any remediation program in the context of students' misconception. Most frequently used techniques for extracting students' ideas are, clinical interview, 2-tier diagnostic test, concept inventories, concept maps and multiple choice tests which incorporate common misconceptions as item distracters. Other approaches have used drawing, sorting and word association tasks and computer simulations. The objective of this study was to develop a CBOT about the subject of "Life Processes" in 9th grade Science curriculum and to argue the efficiency of this test on students' achievement. The researcher developed four Concept-Based Objective Tests (CBOTs).

The CBOTs development procedure had three general steps: defining the content boundaries of the test, collecting information on students' misconceptions, and instrument development. Misconception data were collected from open-ended questionnaire, interviews and content analysis. The data were used to develop CBOTs multiple-choice items.

The researcher decided to use multiple choice questions with four alternatives. The stem of each of the multiple choice items was describing the concepts, structures, processes, or functions. A correct answer

and three distracters were provided for each of the items. After each question, space was provided to write the reason for selection of their answer.

The results of the study showed that the CBOTs were effective on determining the students' misconceptions and also it might be used as an alternative to the traditional multiple choice test for of assessment and evaluation of students' misconceptions. Through these CBOTs, not only the students' understanding of the life processes was revealed but misconceptions among the students were also detected.

Dr. Narendra D. Deshmukh, Homi Bhabha Centre for Science Education, TIFR, V. N. Purao Road, Mankhund, Mumbai 400088, India

Analysis of Biology Education in Japanese School in Terms of Its Priority among Science Subjects

Kseniya Fomichova, Taku Misonou
University of Yamanashi, Japan

The 21st century is referred to as “the era of biology.” However, it is unclear if the 21st century world wide education systems are prepared for this challenge in terms of their structure and contents – if they place an appropriate emphasis on “Biology” in contrast to other science subjects. This research aims to analyze characteristics and factors defining a relative prioritization of physics and chemistry compared to biology in Japanese schools, and to learn if such prioritization is accomplished internationally. Analysis is based on the results of international educational comparison, international and Japanese publications and interviews with stakeholders of Japanese school education.

While PISA shows higher rate of interest of Japanese pupils to biology than to other scientific fields and this subject is chosen by a prevailing number of them, biology is considered to be more suitable for low performers in science. Also, it is supposed to be more appropriate for the mentality of girls than that of boys. International research has discovered that life and environmental science topics are taught to lesser percent of pupils than topics in physics and chemistry in Japan compared to majority of participating countries. OECD indicates that scientific literacy of Japanese population is lower in biological concepts. Some of

these trends are not specific on a cultural basis, but can be found in other countries. A number of them (German, Australian, etc.) experienced and over came periods of undervaluation of biology at different stages of development. Other systems of education, for example, that of Ukraine and Russian Federation presently show certain tendencies similar to those of Japan.

However, new national curriculum standards for school science were introduced from April 2012 in Japan. They aim to end the “minimum curriculum age” and may promote the sustainable development of this subject.

Ms. Kseniya Fomichova, Natural, Biotic and Social Environmental Engineering, Interdisciplinary Graduate School of Medicine and Engineering, University of Yamanashi, Japan

The Role of Bioscience Training Partnerships in Expanding Employment Opportunities for Bangkok's Urban Community Case Study

Churdchai Cheowtirakul
Assumption University, Thailand

This project was initiated during the history worst flooded of Bangkok and vicinity during 2011. The flood cause the massive migrate of many valuable industries from Thailand to countries such as Vietnam, Laos, Cambodia and Myanmar. This study addresses a gap in the providing of golden opportunities in regarding the extent to which public workforce and volunteer in utilizing a biological knowledge-intensive expand employment for a nontraditional, i.e., less-educated or displaced workforce. Specifically, it investigated the problem of knowledge generate institute to help to provided useful education to the communities for mitigation of the disaster phenomenon in Bangkok and urban area. The study also examined the mechanisms through which educators influence volunteer and hiring practices, focusing on strategies to encourage employers' active engagement in the partnership and to facilitate extensive collaboration among key partners. To evaluate impact on community practices, and determine the factors associated with successful (or unsuccessful) program intervention in the sample phenomenon.

We conducted intensive training and publicity to make

the public understand of the basic biological concept that can help to solve the problem in need. The programs have succeeded in ensure of the concept and knowledge we training future technicians to meet the needs of urban communities; and in negotiating skills-based, as opposed to credential-based application. Finally, the evidence shows that partnership efforts produce biological training, and services; facilitate the learning necessary to generate program innovation; and establish relationships of trust with communities. Together, such outcomes positively shape the opportunity structure facing community to create jobs base on biological innovation knowledge.

Prof. Churdchai Cheowtirakul, Faculty of Biotechnology, Assumption University, Thailand

Performance Evaluation of School Divisions in Mindanao: Education for All 2015

Gee Marie S. Binag¹, Rec Eguia²

¹*Agusan del Sur State Agriculture and Technology;*

²*University of Southeastern Philippines, Philippines*

The educational process cannot exist without pedagogical resources: study programmes, textbooks, and instructional materials adapted for the different age groups found within the educational system. Here again, the tasks of conceptualization, elaboration, usage, evaluation, review often fall to the universities. Achieving the “Education for All” goals within the framework of the United Nation’s Millennium Development Objectives requires a concerted effort and synergy amongst all societal players, from the individual to the institutional standpoint. Universities do have a major role to play, cooperating closely with public institutions and civil society. The study was conducted to evaluate the technical and productivity performance of schools divisions in Mindanao from period of 2002 – 2010. The profile factor of input and output variables were computed using Descriptive Statistics whereas the Technical and Allocative Efficiency scores and Total Factor Productivity were calculated using DEA with output orientation while the sources of inefficiency of the schools divisions were calculated using the Tobit Regression and the significance of the difference were calculated using SPSS version 16. Based on the findings, among 48 schools divisions in Min-

danao, 10 of these have achieved full technical efficiency level. Meanwhile, 38 schools divisions were inefficient due to lack of factors inputs like the numbers of male and female enrollees, desk, classrooms and MOOE. Thus, these are the determinants in the number of male and female graduates that would affect the performance that resulted to the inefficiency of the schools divisions. Components of the productivity change are due more to technological and total factor productivity changes with efficiency changes providing a gap.

Dr. Gee Marie S. Binag, Agusan del Sur State Agriculture and Technology, Bunawan, Agusan del Sur, Philippines

Exploring Perceptions of Climate Change in Year Eight Secondary School Students in Victoria, Australia

Bianca Weyers, Anne Wallis, Ty Matthews

Deakin University, Australia

Education is an important platform for providing the skills and knowledge required to address environmental problems, such as, the impacts of greenhouse gas emission. Changing climate is a complex issue that is now regarded as one of the world’s most significant environmental challenges. In Australia, however, very few studies have been undertaken to explore student perception about climate change and examination of the Victorian curriculum reveals that climate change is not a focus for any teaching levels from Grade 5 through to Year 10. Understanding student’s knowledge of climate change can provide the basis for curriculum development and ensure students are well informed about an issue that is likely to have broad environmental, social and economic consequences for future generations.

This paper reports on the results of an investigation designed to gather data about student knowledge and perceptions of climate change. Using a combination of qualitative and quantitative methods, Year 8 science students from three regional secondary schools in south-west Victoria, Australia, participated in the research activities. Results showed that students vary in their level of understanding associated with the causes, impacts, and solutions to climate change. They demonstrated low levels of concern and indicated that their

main source of information about climate change was from the media and to a lesser extent from school. Many students were clearly confused and misinformed about many aspects associated with the influence of green house gas emissions and how it is influencing climate.

During the course of the study, it was also found that demographic factors, such as gender and the location of the school did not influence student responses. The most influential factor appeared to be the method used to elicit these responses. This study has shown that the use of triangulated methods can strengthen research associated with research of climate change education. If this small scale study does in fact provide a representative sample of students within Victorian schools, a greater emphasis on climate change is required to ensure that current students can implement and/or understand the need for adaptation to climate change now and in the future.

Dr. Anne M. Wallis, School of Life and Sciences, Deakin University, Warrnambool, Vic. 3290, Australia

Importance of Parasite Diversity in the Understanding of Global Biodiversity

C. H. Diong, Shancui Serene Oh

Nanyang Technology University, Singapore

A substantial proportion of global diversity is represented by diverse groups of organisms, parasites. Parasites are highly adaptable organisms that have been recognized as contemporary probes of biodiversity. Several groups of parasites are disease agents affecting human health, others are of agricultural, epidemiological, and veterinary importance. The diversity of metazoan parasites of wildlife vertebrates is far from complete. Parasitological surveys and parasite inventories are not evenly represented across taxa. Increasingly however, parasites are included in biodiversity surveys and to study parasite diversity. A number of determinants of parasite diversity seems to have evolved independently in different host-parasite systems. Measures used to study parasite species richness patterns are usually investigated at the infracommunity and component community levels. Parasite diversity provides insight into the structuring of ecosystems, zoogeography, impacts on local biodiversity

by non-indigenous host-transported parasites into naturalized habitats, colonization-extinction dynamics, and importantly, biodiversity on a local, regional, and global scale.

C. H. Diong, 2M Jalan Remaja, Singapore 636671

Educational Implication for Mangrove Rehabilitation in Barangay Catadman Cuyo, Palawan

Victor Magbanua, Marjorie Espanola, Agnes Acero

Palawan State University, Philippines

Barangay Catadman is one kilometer away north of the town of Cuyo. The island town is 156 nautical miles away from Puerto Princesa City. In the past, the residents of Cuyo Island found the place as the source of marine resources which served as their major source of livelihood, but due to climate change and anthropogenic activities, it resulted to the diminished mangroves in the areas. The PSU-Cuyo through its Extension Services had spearheaded a project to rehabilitate the mangroves. Hundreds of mangroves had been planted in the place. A database of some marine resources in the area is also being established for future reference. A program was already started which shows a way of mitigating the effect of climate change by educating the residents of the importance of a well-preserved environment, demonstrate the proper care and rehabilitation of the diminished natural resources. The reasons for the diminishing marine resources are illegal fishing, traditional way of fishing using *lamnek*, *lagtang*, *makaslaand sarap*, cutting mangrove to be used as decoration, firewood, charcoal, stock house building, for the construction of their *ku-boor nipa* hut and *kaingin*. Adding to the degradation of the area observed later was due to the unpredictable climate changes. Much later the use of commercially bought poison for live fish catching had aggravated the degradation of the marine resource. From the views of the residents and observations by the faculty and students, the reasons for the mangrove preservation and conservation are protection from extreme heat, strong winds, large waves, typhoons, windbreaks for the whole shoreline of the community; protection from soil,

beach and sand erosions; breeding and nursery area for marine organisms and shelter for inland animals like reptiles, insects and birds.

Prof. Victor Magbanua, College of Community Resources and Development, Palawan State University, Cuyo Campus, Palawan, Philippines

Implementations of the Lesson Using Diatom Project: Web-based Educational Aids in Seven Language Areas of Asian Countries

Shigeki Mayama¹, K. Katoh², H. Omori², S. Seino³, H. Osaki⁴, M. Julius⁵, J. H. Lee⁶, C. Cheong⁶, E. A. Lobo⁷, A. Witkowski⁸, R. Srivibool⁹, P. Muangphra¹⁰, R. Jahn¹¹, M. Kulikovskiy¹², P. B. Hamilton¹³, Y. Gao¹⁴, L. Ector¹⁵, T. Soeprbowati¹⁶, K. Balasubramanian¹⁷, B. Alakananda¹⁷, S. Guruprasad¹⁷, E. A. Barlaan¹⁸, C. N. Solak¹⁹

¹Tokyo Gakugei University, ²Tokyo University, ³Kyushu University, ⁴Stream Graph, Japan; ⁵St. Cloud State

University, USA; ⁶Daegu University, Korea; ⁷University Santa Cruz do Sul, Brazil; ⁸University Szczecin, Poland;

⁹Brapha University, ¹⁰Silpakorn University, Thailand;

¹¹Bot. Garten Bot. Museum Berlin-Dahlem, Germany;

¹²Institute for Biology of Inland Waters, Russian Academy of Science, Russia; ¹³Canadian Museum of Nature,

Canada; ¹⁴Xiamen University, China; ¹⁵Public Research Centre-Gabriel Lippmann, Ruxenburg; ¹⁶Diponegoro

University, Indonesia; ¹⁷Gubbi Labs, India; ¹⁸UP-Diliman, Philippines; ¹⁹Dumlupinar University, Turkey

Diatom Project is an international Web-based educational system for promoting awareness to the nations about river environment*. In the past three years, we have developed it in 17 languages with four modules: SimRiver – a Web-based program simulator that uses diatoms to enhance the understanding of the relationship between human activity and water quality; streaming videos – introducing basic knowledge of diatoms; visual tools – informing circumstances of various water environments in both different areas and time frames; and reporting system – for feedback from learners.

Effectiveness of the system was assessed in classes using six languages in five countries. Before and after the classroom activity, students described answer for the same question: What thought do you have about river today? The descriptions were treated with text mining software followed by correspondence analysis (CA). In comparison of the pre-and post-tests, Japan

and Korea showed similar shift of student groups in which their awareness was oriented toward improvement of riverine environment. In Indonesia and two Indian language (Kannada and Marathi) areas, they were rather oriented to SimRiver program and diatom itself. Thai students showed intermediate shift between eastern and southern Asian countries. The difference of the group positions in CA may be caused by different learners' awareness about river environment for sustainable development and will allow us to invite people from any country to join the project. Very recently we had a class activity using the Diatom Project in the Philippines. We will also show the result of this analysis in comparison with those of other Asian language areas.

* <http://www.u-gakugei.ac.jp/~diatom/>

Prof. Shigeki Mayama, Tokyo Gakugei University, Tokyo, Japan

Comparative Assessment in Using Textual Graphical and Traditional Instructional Material in Teaching Biology

Christine May Torres¹, Ana P. Centeno²
¹Columban College Inc., ²Colegio de Calumpit, Philippines

The Philippine education, particularly the basic education, is now faced with a great challenge upon implementing the K-12 scheme of education. One of the challenges concerning this is the selection of appropriate instructional materials. Instructional materials play very important aspect in the teaching and learning. In fact, this is one of the many factors that are blamed for the poor quality of education in the country. One of the predicaments is addressing the issue of the instructional materials appropriate and suitable with the type of learners. Sufficient theories and literatures are available stating the advantages and disadvantages of using instructional materials presented in heavy text, graphical and traditional means. Therefore, as a prelude, this study's main concern was about assessing the significant effects of text, graphical and traditional presentation of instructional material. In the conduct of the experimental research, the tool used to determine the significant difference on the performance of students in particular topics in biology intervened by

heavy text module, graphical module and the traditional instructional material. As a result, in its finding, it shows that there is no significant difference between students' performance intervened with book and those that intervened with text. Also, there is no significant difference on the performance to those intervened with book and those with graphical material. However, there is a significant difference between those students intervened with graph and text. This draws the conclusion that students learn and perform more if they will be using more graphic material. From this study, it is further recommended to construct instructional materials with more graphics. Also, apply the graphical concept to other topics in Biology as well with other subjects. Lastly, the study may also be utilized in other subject matter for validation.

Opinions of Japanese University Students about Issues of Bioethics: The Case of Gene Diagnosis and Amniotic Fluid Examination

Kunio Umeno¹, Junko Iwama², Shizuo Matsubara³
¹Japan University of Economics, ²Kawasaki City College of Nursing, ³Toin University of Yokohama, Japan

We investigated Japanese university students' opinions on "gene diagnosis" and "amniotic fluid examination" as examples of bioethical issues, which are important component of biology education, by questionnaire in July 2012. The number of respondents was 322 from three universities. The questions were "Would you want to have a gene diagnosis, if this examination were available for any applicant?" and "Would you want to have an amniotic fluid examination for yourself or your spouse, after you married and you or your spouse were pregnant?" Respondents chose one answer from four alternatives: (1) I strongly hope so, (2) I hope so, (3) I do not hope so, and (4) I strongly do not hope so. According to the collection of all responses, the most common response was alternative (1) and the second was alternative (2) for both questions. Total rates of selection of positive answers, namely alternatives (1) plus (2), were 83.7% for "gene diagnosis," and 73.8% for "amniotic fluid examination." There were little gender differences in the tendencies of selecting alter-

natives. As to "gene diagnosis," the selection rate of alternatives (1) and (2) were 45.3% and 36.7% respectively for male students, 39.4% and 46.8% respectively for female students. As to "amniotic fluid examination," total selection rates of positive alternatives, (1) plus (2), for male and female students were 76.2% and 70.2%, respectively. As shown in the above data, there was a tendency that female students have relatively more negative opinions than male students for having "gene diagnosis" and "amniotic fluid examination." However, the cause of this tendency may not gender difference, but might be that the majority of female student were nursing majors.

Opinions of Japanese University Students about Issues of Bioethics: The Case of "Organ Transplantation and Brain Death" and "Preimplantation Genetic Diagnosis"

Junko Iwama¹, Shizuo Matsubara², Kunio Umeno³
¹Kawasaki City College of Nursing, ²Toin University of Yokohama, ³Japan University of Economics, Japan

Bioethics relates to every aspect of life in nature and social environments. In recent years, the necessity of guidance about the "preciousness of life" has been proposed by many researchers for science education. However, our research has revealed that in Japan, implementation rate of bioethics education in lower secondary school was about 20%, and in elementary schools and upper secondary school, it was only about 10% (Iwama *et al.*, 2012). We investigated Japanese university students' opinions on "organ transplantation and brain death" and "preimplantation genetic diagnosis" in July 2012. The results of the study are as follows: 1) Firstly, on "organ transplantation and brain death," 65% of the students (91 of 140) approved "conditional promotion," that is, "I accept both organ transportation and brain death, and I hope artificial organs will be developed." The number of "promotion" students was 17%. So the great majority of students (82%) accepted the promotion of organ transplantation. 2) Secondly, on "preimplantation genetic diagnosis" in order to treat the first child who is suffering from "fanconi anemia," 37% of the students (52 of 140) approved "conditional restriction," that is, "We should not

request the second young child who has no will to become the means of the treatment.” The number of students answering “restriction” or “negation” was 27%. Thus the majority of students (64%) had negative opinion. Japanese students usually study about the issues of bioethics at classes such as “Integrated Studies,” “Ethics” and “Contemporary Society” in upper secondary school. However, bioethics relates closely to the body and life of living things, so it also should be taught in school science education.

Alternative Conceptions of Human Cardiovascular System Concepts Among BS Nursing Students in Zamboanga City

Obra, Mario

*Western Mindanao State University and Philippines
Ateneo de Zamboanga University, Philippines*

This study was conducted to elicit the alternative conceptions of human cardio-vascular system (HCVS) concepts among BS Nursing students in Zamboanga City. Specifically, the study aimed to identify the specific alternative conceptions of students on heart structure and functions; nature and components of blood; blood functions; blood vessel structure and functions; blood circulation, HCVS general function; HCVS disorders. The respondents of the study were 312 BS Nursing students coming from the Ateneo de Zamboanga University, Western Mindanao State University and Univeridad de Zamboanga. A random multi-stage sampling design was used to select the actual respondents from each university. Students’ alternative conceptions on essential content areas of HCVS were identified through the analysis of items and choices of the respondents on the researcher prepared two- tiered diagnostic test. Individual responses and group output of the respondents were noted and identified. Results showed that BS Nursing students in Zamboanga City continued to have alternative conceptions on HCVS concepts even after formal instruction. The students had the highest percentage of alternative conceptions on blood components (65%); HCVS general function (52%); understanding hypertension (46%) and nature of blood (38%). The least percentage of alternative conceptions was detected on blood circula-

tion (3%), heart structure (5%) and blood functions. Hence, these findings support the claim that HCVS is a topic in Biology where students often hold considerable number of alternative conceptions.

Prof. Mario Obra, Western Mindanao State University, College of Science and Mathematics, Zamboanga City, Philippines

Using TED As Supplementary Materials for Teaching Biology

**Kyoungho Kim¹, Kew-Cheol Shim²,
Jung-min Kim³, Nam-Il Kim⁴**

¹Gongju National University of Education, ²Kongju National University, ³Environmental Ecological Education Institute, ⁴Chuncheon National University of Education, South Korea

It is mainly science textbooks that provide information to students. However, textbooks fail to cover the latest scientific developments, as their information is at least 10 years old. The most recent scientific progress is often introduced through various materials from simple black-and-white photos to more sophisticated videos. One of the most up-to-date materials is TED videos. TED is open to anyone with access to the Internet via computers or smart phones. Speakers of diverse backgrounds share their unique ideas through a 5 to 20-minute lecture on various themes and videos of those lectures are updated every week. This presentation aims at introducing some of the TED videos that could be used for teaching biology: functions of mushrooms, honeybee breeding, colour recognition of a monochromat, life in the deep oceans, and medical use of archeological discoveries.

Prof. Kyoungho Kim, Department of Science Education, Gongju National University of Education, Gongju, Chungnam, South Korea

Analysis of Japanese Science Curriculum for Connecting to Our Daily Life: Crosscurriculum with Health Education About Human Bodies

Sato, Takayuki

Hirosaki University, Japan

Science education has necessity for connecting the learning contents to daily life in Japan. Therefore, we established purpose of this study which connects learn-

ing human bodies to daily life with using health education. So, we compared Japanese science curriculum with health education curriculum. The results of analysis of “Japanese Course of Study,” learning contents about human bodies have been in science curriculum all periods after World War II. The results of analysis of science textbooks which is used in compulsory education, bone structures, muscles, embryogeny, respiration, digestion, blood circulation and principal internal organs are learned at the latter half grades of elementary schools. On the other side, at lower secondary schools, nervous systems, reproduction, and advanced contents of elementary school level are learned. We analyzed learning contents which were described in health education textbooks. There are many learning contents to understand health on daily life at elementary schools. The learning contents of health education at lower secondary schools are to understand health concretely with viewpoints of structure and function of human bodies. Therefore, when the contents of health education are introduced in science learning, science curriculum can connect to daily life strongly. If teachers use the contents of health education at introduction of science learning activities, students become to have motivation to learn science by feeling affinity to science. If teachers use them at the last of science learning activities, teachers become to be able to emphasize the connection of science learning and daily life. Furthermore, there are many modernistic problems about human health which are explained scientifically on health education textbooks. If these problems are applied to science learning activities, students have opportunities that they use basic knowledge of science that they have learned. Then, we think that students’ desire to learn science can be improved.

<Poster presentations>

Assessing the Potential for Transformative Learning in a Field Biology Course

Chuckie Fer A. Calsado, Dawn T. Crisolago, Justin Ray M. Guce

Biology Unit, Philippine Science High School – Main Campus, Philippines

The Field Biology elective is a three-week program

offered to incoming senior students of the PSHS-Main Campus. Participants undergo rigorous on-campus training, team-building, and orientation activities prior to being deployed to a variety of ecosystems in Luzon where they employ field techniques to collect and analyze data on ecological interactions and human impacts on the environment. Its informal, experiential, collaborative, and immersive nature makes it a good venue for effecting transformative learning. Quantitative data in the form of scores on Frederick’s Cognitive Reflection Test (2005) and Strathman Consideration of Future Consequences Scale (1994) showed that a majority of the course participants focused more on the potential future outcomes of their current behavior, a trait associated with increased environmental awareness. Qualitative data from interviews and reflection papers showed increased interest in the biology as a career choice, increased awareness of environmental issues and modification of unsustainable behavior, and development of life skills. Learning experiences that have contributed to these shifts in thoughts, feelings, and behaviors are also identified.

Development and Practice of Pre-service Teacher training Program for Elementary Science: A Plan for Teachers’ Competency in Biological Observation and Experiment in the New Japanese Course of Study

Heiwa Muko, Masahiro Hizume, Hiroshi Leyama
Ehime University, Japan

Since 2011, the new Japanese Course of Study has been implemented. One salient change in the Course of Study was the increase in the number of contact hours for elementary science, from 350 hours to 450 hours. New observation activities and experiments have been also brought into the new elementary science. The purpose of this study was to develop a special pre-service teacher training program for elementary science teaching guided by the new Course of Study, and to investigate the effects of the program to prospective elementary teachers. The training program was conducted for 26 university hours filled with observation activities and experiments, which covered most of the science contents in the new Course of

Study. A total of 800 prospective teachers from various departments and/or colleges of Ehime University participated for the academic year 2011-2012. It can be implied in the results that the participants acquired basic scientific knowledge and competency in observation activities and experiments through the program. Although the program is non-credit, prospective teachers who actively participated in at least 70% of the list of activities were given certificate of completion. The success of the program is deemed important in improving and ensuring the high quality of pre-service teacher training program for elementary science teaching.

Dr. Heiwa Muko, Department of Education, Ehime University, Japan

Do Bell Peppers Photosynthesize? ~ Inquiry into Photosynthesis ~

Teiko Nakamichi

Tokyo Institute of Biology Education, Japan

In elementary and lower secondary school, students learn about photosynthesis: leaves of a plant catch sunlight and make nutrition such as starch. How do students conceptualize photosynthesis occurring in different parts of the plant? I surveyed secondary school and university students. The key question I asked the students was, "Does the fruit of red bell peppers and green bell peppers photosynthesize?" Thirty-seven percent of the students answered that red bell peppers do photosynthesize, while 49% answered that green bell peppers do not photosynthesize. Some fruits contain chloroplasts, like leaves do. Students will be better able to understand that flowers evolved from leaves by inquiring more about photosynthesis in the fruit of bell peppers. Furthermore, by including peapods in the inquiry activity and by observing stomata in the epidermis of peapods students can recognize the commonality of leaves and fruits. The experimental method was as follows: (1) Prepare the materials such as bell peppers of different colors, peapods, and spinach leaves. Cut each of them into an appropriate size so as to keep each one at the middle of a test tube. (2) Put 1 ml of pH indicator in the bottom of the test tube (In this step, I used a mixed solution of thymol blue and phenol red as a pH indicator). (3)

Put materials in the test tube avoiding contact with the pH indicator solution, and make the test tube airtight with a rubber stopper. Prepare a test tube with only the pH indicator in it as a control. (4) Prepare another set of test tubes in the same way, and wrap them in aluminum foil to block light. (5) After placing all test tubes under a light for a while, examine the color of the pH indicator in each test tube.

Mrs. Teiko Nakamichi, e-mail: teiko-n@nifty.com

Elementary School Teachers' Beliefs about Education for Sustainable Development

HyeongCheol Lee, PyoungKil Yoo, JinHo Bae

Busan National University of Education, South Korea

The purpose of this study was to investigate elementary school teachers' belief about education for sustainable development (ESD) in South Korea. Subjects were 75 (male 25, female 50) elementary school teachers. Questionnaires, developed on the basis of Yang *et al.* (2010) which was for measuring of the beliefs of secondary school teachers, consisted of values of ESD (respect and care for the community of life, ecological integrity, social and economic justice, democracy, non-violence and peace) and teaching beliefs of ESD (relevance to daily life, students' need in the future, integrated teaching). The results of this study were as follows: Elementary school teachers generally showed positive beliefs about values of ESD and teaching beliefs related to curriculum content, pedagogy, and learning and teaching beliefs of ESD. But they showed negative beliefs about social and economic justice comparatively.

Prof. HyeongCheol Lee, Department of Science Education, Busan National University of Education, Busan 611-736, South Korea

Development of Environmental and Experiential Program for Understanding the System and Environment of Local River

Kiyoyuki Ohshika¹, Kazunari Saguwara²

¹Aichi University of Education, ²Foundation of River & Waterland Environment Management, Japan

In the new Japanese Curriculum, the integration of real life experiences, utilization of scientific museum, and

the use of local resources in teaching science were given emphasis as strategies in science teaching. The idea was applied to the use of following three rivers: Kiso River, Nagara River, and Ibi River located in Chubu area at the center of Japan. The area is rich in natural resources with high diversities of living things around. The place could be served as a good study site in which different topics and concepts in environmental science can be demonstrated in different year level of education, such as functions of river for grade 5, the living things and their environment for grade 6, and physicochemical survey of water quality of the river for the junior high school level. In this research, the environmental educational program was developed with a theme intended for local rivers for the students to learn local natural environment and living things, and to practice formal and informal educational opportunities. This program has three main themes based on the characteristics of Kiso River: the natural environment, commercial use, and the tradition and culture of the local people. In this research, we focused on the natural environmental section. Using this program, students experienced real life situation or living things in local rivers, and teachers could teach about the river without having special skills. Because students are able to learn about the local rivers experientially, they will continue to learn by themselves and develop their own understandings in the future.

Dr. Kiyoyuki Ohshika, Department of Science Education, Aichi University of Education, Japan

Origami Bird: A Teaching Material Linking Mutation, Natural Selection, and Speciation

Takahiro Yamanoi^{1,2,3}, Watal M. Iwasaki⁴, Masaharu Takemura², Vivian Tolentino³

¹Hakuoh Ashikaga Senior High School, ²Tokyo University of Science, ³the University of Tokyo, ⁴Tohoku University, Japan

We improved the origami bird protocol (Westerling, 1992), a teaching material for natural selection, by incorporating Gametes mutation box that links gene alteration to changes in phenotype and Origami bird simulator that links natural selection and speciation. A comparison between the pre-and post-test results suggested that this protocol successfully enhanced

Japanese high school students' understanding of linkage between mutation, natural selection and speciation, and removed misconceptions about topics such as Lamarckism and orthogenesis. Our improved protocol could be used widely because, in many countries, teaching materials linking microevolution and macroevolution are lacking (Catrely, 2006) and foreign students also have teleological conception about evolution (Bardapurkar, 2008).

Prof. Takahiro Yamanoi, Hakuoh Ashikaga Senior High School, Tadaki-cho 1067, Ashikaga, Tochigi 329-4214, Japan

Development of the Vegetable-based Methods for Learning Molecular Systematology in Japanese High School Biology Course

Tomoko Kazama, Masaharu Takemura

Tokyo University of Science, Japan

Over the past several decades, tremendous growth has occurred in our understanding of genetic phenomena and the intricate and complicated mechanisms that mediate genetic effects. During study in secondary schools in Japan, modern genetics is an important area for high school students. Now we are developing new teaching materials for learning molecular systematology, which includes processes necessary for producing a molecular phylogenetic tree by the students themselves. Because vegetables are regarded as familiar for students, we are trying to use them as new teaching materials. We introduced the hypothesis that "the edible portion of vegetables and the form of a flower reflect the systems of these vegetables" to this teaching material. We established methods of making phylogenetic trees using sequences of *rbcl* genes of some vegetables. Vegetable-based methods are thought to incorporate "artificial selection," and are expected to include effective educational points related to species and scientific names. Using these teaching materials, students are thought to be able to learn about molecular evolution, molecular phylogenetic trees, molecular clocks, scientific names, and classification systems of living things. Herein, we present the results obtained by application of these teaching materials for Japanese school students.

Ms. Tomoko Kazama, Graduate School of Mathematics and

Science Education, Tokyo University of Science, Kagurazaka 1-3, Shinjuku, Tokyo 162-8601, Japan

Development of a Simulation Activity for Learning Evolution

Tetsuya Asano, Kiyoyuki Ohshika

Aichi University of Education, Japan

In the new Course of Study of Japan, it is to learn biology from a perspective of regularity and biodiversity at the high school level. "Evolution" is one of the important topics to learn in biology. However, it is difficult to observe directly the process of evolution and to conduct experiments. In addition, it is difficult to develop teaching materials for the topic of evolution. In this research, we developed a simulation activity on natural selection (based on the activity of Burton *et al.*, 2009), and observed the effectiveness of this activity in the teaching and learning process. In the activity, students use table utensils such as spoon or fork. Students observe the change in the number of utensils during the conduct of this activity. The activity was tried out to the undergraduate science major students to find out its effectiveness in developing the concepts in evolution. Before and after the conduct of the activity, the students were given questionnaire to assess their understanding about natural selection as well as their views about the activity. From the results of analysis on students' responses, many students considered the activity was very interesting and easy. Furthermore, they could understand the process of evolution in this activity. Therefore, we recommend this activity as an effective material to teach natural selection in the topic evolution in high school biology.

Mr. Tetsuya Asano, Graduate School of Education, Aichi University of Education, Japan

Development of Teaching Material Using Owl Pellet of the Zoo

Shiho Chiga, Kiyoyuki Ohshika

Aichi University of Education, Japan

Raptors are known for vomiting pellets that contain undigested matter such as bones or hairs. Pellets are used as a general method of investigating the feeding habits of raptors. Owl pellets are used as a teaching

material for food webs in the United States. Ohshika *et al.* (2010) carried out a lesson with owl pellets as a teaching material imported from the United States; it was suggested that pellets are effective as a teaching material for food webs at junior high schools. However, pellets are not familiar for Japanese children and it is a problem for the teachers to identify the contents of the pellets. Under the new Japanese Course of Study, real life experience, utilization of science museum, and application of local resources are given emphasis. In this research, teaching material of owl pellets from a local zoo in Japan was developed. Owl pellets were collected from nine zoos in Japan, and subjected to content analysis. As a result, owls in many zoos consume mainly two types of animals, either mouse or chicken. Therefore, the "bone charts" of a mouse and a chicken were developed. Then, the "bone charts" and the pellets were introduced in science classes at the 8th grade and 9th grade. Results show that many students could examine pellets positively and the students' ideas about the owl being herbivores were changed into carnivorous animals. Furthermore, they are more interested in the feeding habits of other animals and morphological features of the skeleton of the mammals. Therefore, we believe that owl pellets are effective teaching material to understand the feeding habits of the owls.

Ms. Shiho Chiga, Graduate School of Education, Aichi University of Education, Japan

Effect of Learning DNA and Gene Using an Experiment of DNA Extraction at Junior High School

Toru Higuchi, Kiyoyuki Ohshika

Aichi University of Education, Japan

In the new Japanese Course of Study, learning the concepts of DNA and gene through observations and experiments and understanding the regularity of the organisms through the study of the nature and structure of DNA are emphasized. DNA extraction is now included in the new junior high school science textbooks in Japan. However, this experiment has hardly been done in junior high school because of some problems about protocols and effects. In this research, the best protocol for conducting the experiment on DNA ex-

traction was tried out in 9th grade students. Pre- and post-questionnaires were administered and analyzed to determine the effectiveness of the activity. It was revealed by the pre-questionnaire that most of the students had known the term DNA through TV programs but many of the students thought that plants are not organisms and do not have DNA. From the results of pre-questionnaire, it is important for students to experience DNA extraction from both animal and plant. The “cod roe” (an animal material) and “banana” (a plant material) were used for the experiment and students could extract DNA from these materials. Students’ responses in the post-questionnaire showed that they realized plants are organisms and all organisms have DNA. Many students could realize the correct concept about DNA through this activity. We believe that misconceptions of junior high school students regarding DNA and genes will be corrected through this activity.

Mr. Toru Higuchi, Graduate School of Education, Aichi University of Education, Japan

Science High School Students’ Conceptions about Climate Change

Arnold D. Pitpitunge

Philippine Science High School – Central Luzon Campus, Philippines

There were no studies that focused on students’ correct conceptions (extent of knowledge and comprehension), misconceptions, and alternative conceptions on the basic concepts, causes, effects, mitigations of, and adaptations to climate change. This study determined the science high school students’ conceptions on the aspects of climate change. It involved randomly selected 122 science high school students who took the researcher-made multiple choice climate change test. In each item, the students encircled their choice, provided brief explanation about their answer, and determined the source of information about the concept. Their answers were coded to determine their climate change conceptions. Students hold correct conceptions on the observable phenomenon that indicates climate change, product that produce greenhouse gases, direct effect of climate change on freshwater, activity to participate in for climate change mitigation, and strat-

egy to cope with the impact of climate change on biodiversity. However, misconceptions on relationship of climate change and weather, factors that change the Earth’s climate change, effects of climate change on coastal areas, products to buy to support carbon sequestration, and strategy to cope with the impact of extreme weather conditions on agriculture were revealed. Furthermore, more alternative conceptions (ideas, views, beliefs about climate change that are not consistent with scientifically accepted views but are not considered to be errors or incorrect) on relationship of climate change and health problems and strategy to cope with freshwater stress than the other concepts on climate change were also noted. Male respondents have more correct conceptions than female respondents who have more misconceptions on climate change mitigations. Second year respondents have more correct conceptions; the first year respondents have more alternative conceptions. Information from home/community decreases the incorrect views and alternative conceptions of students on the basic concepts of climate change. Media increase misconceptions but decrease the alternative conceptions on the effect of climate change.

Prof. Arnold D. Pitpitunge, Philippine Science High School – Central Luzon Campus, Clark Freeport Zone, Pampanga, Philippines

Small Group Instructional Diagnosis (SGID): A Reinforcement Strategy

Sheroda S. Montanez, Jovelyn L. Orquina

Father Saturnino Urios University, Philippines

This mixed research study focuses on the Small Group Instructional Diagnosis (SGID) to improve academic performances of the students. The chosen respondents were the classes of Microbiology and Humanities both with two sections. The respondents were purposely chosen to compare whether the SGIS Process can really help improve the performances of the students. The simple random-lottery sampling was used to level off the respondents for the Non-SGID groups. To determine the differences between the academic performance of the same group – Class A1 (Microbiology) and Class A2 (Humanities), the paired *t*-test was used. To determine the differences between the aca-

ademic performance of the two groups, SGID and Non-SGID, the *t*-test was employed. Results revealed different approaches that the teachers used during the class discussion that included group activities, individual activities, class participation and handouts. According to the students, these approaches help them improve their learning. If they stated different approaches that enhanced their learning, they also have distracters that detract their learning that includes “noise.” Majority of them voted noise as one of the most known distracters that really detracts them from learning. The grades of the students from the four classes, Class A1, Class A2, Class B1 and Class B2, increased, but the SGID groups had higher improvement compared to the grades of Non-SGID groups. It was also shown that there was a significant improvement on the academic performance of the students who were administered with SGID. Also there was a significant difference between the groups of SGID and Non-SGID. Thus, SGID is significant in improving student’s good performance.

Elementary School Teachers’ Knowledge and Attitudes Towards Renewable Energy Sources

Keum-Hyun So, Yong-Gwon Kim, Seok Hee Lee
Busan National University of Education, South Korea

The purpose of this study was to investigate knowledge and attitudes of elementary school teachers in South Korea towards Renewable Energy Sources (RES). Subjects were 74 (25 males and 49 females) elementary school teachers. Questionnaires, developed on the basis of Liarakou *et al.* (2009) to measure the knowledge and attitudes of elementary school teachers consisted of socio-demographic characteristics of teachers, general knowledge of RES and sustainable development, general attitudes towards RES, attitudes towards wind and other solar farms, domestic uses of RES, teaching about RES. The results of this study were as follows: Elementary school teachers (1) generally showed affirmative perceptions of RES to sustainable development; (2) expected RES to be used as main future energy sources. But in-service training, sustainable educational material, lesson plans, and ex-

tracurricular educational programs in order to deal with RES are still needed by the elementary school teachers in South Korea.

Prof. Keum-Hyun So, Department of Science Education, Busan National University of Education, Busan 611-736, South Korea

Using Scientific Research-based Monitoring Model for Scientifically Gifted Students

Kew-Cheol Shim¹, Ji-Hyon Kil²

¹Kongju National University, ²National Institute of Environmental Research, South Korea

The purpose of this study was to examine educational implications of the scientific research based mentoring model (SRMM) for scientifically gifted students, which was proposed by Shim and Kim (2005) and Shim *et al.* (2009). The SRMM consists of six phases, *i.e.*, introducing, exploring, planning, performing, verifying and concluding, to facilitate scientific inquiry of gifted students in the process of experimental performance. In addition, they could be provided with concrete experimental experiences through laboratory practices, and challenging tasks and feedback at the requisite stage. In this paper, the model was applied to biological research theme, which was the auto-inhibition and inter-inhibition of germination and seedling growth during early stages of plant growth, and educational potentials were examined in Science Education Institute for the Gifted, adjacent to University, Korea.

Prof. Kew-Cheol Shim, Department of Biology Education, Kongju National University, Gongju, Chungnam, South Korea

The Effects of Science, Technology, Engineering, Art, Mathematics (STEAM) Programs in Biology and Sports Science

**Juneuy Hong, Hwagun Byun, Byungyong Lee,
Kyungjun Ahn, Jinhee Park, Hyokwan Dong,
Jaeho Sim**

Seowon University, Korea

Now, there is paradigm shift to the society of creative intelligence with emotion. In the future, it will be important not only the creative personality and emotion but also artistic merit. So we adapt Science, Technology, Engineering, Art, Mathematics (STEAM)

education as a convergence education. The purposes of this study were to analyze students' perceptions about convergence study and to find out the effectiveness of STEAM programs. For this study, we had developed STEAM programs on bio and sports sciences. Fourth graders, 6th graders, 8th graders and 10th graders were participated. The results showed that the STEAM programs were very interesting for students and effective to promoting creativity in science.

Prof. Juneuy Hong, Department of Biology Education, Seowon University, Korea

Various Laboratory Experiences Facilitate Gifted High School Students to Write Scientific Research Articles

**Ji-Hyon Kil¹⁾, Kew-Cheol Shim²⁾, Kyoungho Kim³⁾,
Jung-Min Kim⁴⁾**

*^{1)National Institute of Environmental Research,}
^{2)Kongju National University,}^{3)Gongju National}
University of Education, *^{4)Environmentl Ecological}*
*Education Institute, South Korea**

The purpose of this study was to analyze articles of scientifically gifted high school students, which were published in Korean Journal of Young Scientist. Scientists reviewed the papers and most of the research papers were results of scientific research-based mentoring programs. High school students performed the scientific mentoring programs with the assistance from university professors or institute scientists. Their research themes were related to mathematics, physics, chemistry, biological sciences and earth sciences. Articles related to biological sciences reflect experimental methods and tools including DNA and protein works. The results implied that various laboratory experiences motivated and facilitated them to understand biological properties, and writing an article should be useful to facilitate scientific ability of scientifically gifted students, and to understand the nature of scientific researches.

Prof. Kyoungho Kim, Department of Science Education, Gongju National University of Education, Gongju, Chungnam, South Korea

Integrating Multiple Methods in teaching Biology to Improve Achievement of Students with Varied Learning Styles

Erwin P. Elazegui

Technological University of the Philippines, Philippines

This research investigated the learning styles of students enrolled in general Biology second semester of SY 2011 – 2012 and assess the effectiveness of four different teaching methods introduced into Biology classroom. Cooperative learning, class discussions, concept maps and lecture were integrated into the course to compare students' levels of participation. The study utilized qualitative and quantitative data in an attempt to explore students' conception of selected topics of biology. The Kolbs Learning Style inventory was used to categorize students learning styles and Biology Concept Test was used as research instruments. Results showed that out of 42 students, 24 students were convergers, 9 students were divergers, 4 students were assimilators, and 5 students were accommodators. based on thinking encouraged most of the class participation with a mean score of 4.2. All three methods, however, had a significantly greater level of student involvement than in traditional lecture. The use of different methods caused significant increase in the achievement of biology students with varied learning styles.

Prof. Erwin P. Elazegui, College of Science, Technological University of the Philippines - Manila, Philippines

A Correlational Study on Catechol-O-Methyltransferase (COMT) VAL 158MET Single Nucleotide Polymorphism and Academic Performance of Selected Senior Public High Schools in Manila, Philippines

Janina S. Bautista, Katrina Lara L. Bravo, Luisa Marie S. Cruz, Franz Ysrael R. Del Rosario, Angelo Alexan R. Tanguio, Josefino R. Castillo

University of Santo Tomas, Philippines

Catechol-O-methyltransferase (COMT) is a methylation enzyme that catalyzes the degradation pathway and inactivation of dopamine which has been widely accepted in playing a role in the mediation of human cognition, emotional response, executive cognition and

working memory. The COMT genotypes of student-respondents, determined through DNA isolation, amplification and restriction endonuclease digestion, were correlated with their final grades in school and scores from National Career Assessment Exam (NCAE), an annual assessment program held by the Department of Education to evaluate student's strong points and help them for choosing future career. Results showed that COMT ¹⁵⁸Met/Met homozygous students showed a trend towards poorer performance in mathematics (academic subject) ($p = 0.035$) and in all subtests of the NCAE as compared to students homozygous for ¹⁵⁸Val/Val ($p = 0.036$) or heterozygous for ¹⁵⁸Val/Met ($p = 0.002$). However, the performances of the COMT Met/Met homozygous students in the science and mathematics subtests ($p = 0.001$ and $p = 0.01$, respectively) were significantly poorer than that of the other two groups. The genotypic distribution of the samples in the study revealed 34.2% Met homozygotes, 29.5% Val homozygotes, and 36.2% Val/Met heterozygotes. The low percentage of Filipino students homozygous for COMT ¹⁵⁸Val allele translates into poor performance in the subject areas mentioned above. Numerous educational system amendments have been proposed as they often associate the academic achievement of the students to the effectiveness of the educational scheme. Very little light has been shed on other aspects such as psychological, social and even biological factors that actually affect the academic performance of students.

Prof. Josefino R. Castillo, Department of Biological Sciences, College of Science, University of Santo Tomas, Manila, Philippines

Unraveling the Magnificent Code of Life: Modular Approach in Teaching the Structure and Function of DNA in High School Biology

Shela V. Ruiz, Leah V. Carballo

Central Bicol State University of Agriculture, Philippines

This module generally aimed to present clearer understanding of the structure and function of DNA through activities and discussions primarily designed for second year high school students. The 5 E's of learning was used as a method in the presentation of each lesson in

this module. This includes Engage, Explore, Explain, Elaborate and Evaluate. The module engaged students in practical work activities such as simulation, picture puzzle, model constructions, brainteaser sessions, laboratory investigations, and group output discussion were used. Interaction and collaboration through maximum "minds-on" activity accompanied with "hands-on" and "hearts-on" activities were used as strategies in this module. These activities were done in games and group competition. The discussion of the learning concepts was through teamwork and collaborative learning among students. An output presentation was done every after activity either independently, in pairs, or in groups. The in-depth discussion included in this module provided the teacher a clear understanding on the main concepts to be learned by the students. Finally, this module integrated how the genetic information from parents to offspring is expressed through the central dogma of molecular biology. This would help the teacher and students relate those concepts to the role of the genetic material in the perpetuation of every living organism.

Prof. Shela V. Ruiz, Central Bicol State University of Agriculture, Camarines Sur, Philippines

Two Novel Endemic Philippine Species in *Gynothodes* Blume (Morindeae-Rubiaceae) and Molecular Support on the Reclassification of *Morinda elliptifolia* Merr. and Quisumb

**Joshua Johnedel P. Salvacion¹, Gerson C. Cntreras¹,
Wei Yi D. C. Hung¹, Fernan Macrin C. Ramos¹,
Diana Grace B. Ysaac¹, Jayson G. Chavez²,
Grecebio Jonathan D. Alejandro¹**

¹*University of Santo Tomas, ²Far Eastern University, Philippines*

Recent phylogenetic works on the systematics of Morindeae have proposed new generic delimitations of the tribe and the adoption of a narrow circumscription of the nutraceutical genus *Morinda* known as "noni." The proposed transfer of all lianescent *Morinda* species including the Philippine endemic *M. elliptifolia* to its conglomerate *Gynochthodes* have raised the question whether this taxonomic amendment is supported by molecular dataset. To address this, samples of *M.*

elliptifolia and two *Gynochthodes* cf. specimens were collected in the island of Palawan. *53trnT-F* (cpDNA) sequences were utilized for cladistic analysis. Bayesian inference (BI) of the plastidial data supports the generic transfer proposed by Razafimandimbison & Bremer with strong divergence values (PP = 1.00). The genus is united by marginal hairs along stipules and bracts; axillary, racemose or cymose inflorescences with white and shortly pedunculate flowers; recurved calyx tubes; and corollas with long hairs within the tubes and on the adaxial side of the lobes. Furthermore, the two sampled *Gynochthodes* cf. nestled on the basal polytomy of *Gynochthodes* subclade proving their generic affinity (PP = 1.00). Comparisons between the Malesian *Gynochthodes* and these specimens have shed light to the proposal of two new endemic *Gynochthodes* species. In relation to these taxonomic breakthroughs, implications of Philippine biodiversity, and the industrial and medicinal applications of *Gynochthodes* are presented.

Prof. Jayson G Chavez, Department of Biological Sciences, Institute of Art and Sciences, Far Eastern University, Nicanor Reyes St., Manila, Philippines.

Structural Development and Morphometric Analysis of Male and Female Gametophytes of Nitrogen-absorbing *Canna x generalis* Bailey (Banderang Español)

Emmanuel D. Delocado, Antonio Delfin C. Sumabat IV, Alyssa Jamie W. Gaw, Juan Miguel L. Laurel, Vivian S. Tolentino

Ateneo de Manila University, Philippines

Because of their nitrogen-absorbing ability, canna beds have been used as an ecological remediation strategy for polluted water treatment. In the Philippines, coincidentally, banderang español or canna lily is planted in the road islands of flood prone areas such as España and Marikina. This study aims to trace the ontogeny of *Canna x generalis* Bailey (banderang español) gametophyte and to characterize the development patterns using morphometrics and statistical tools such as repeated multiples ANOVA, *t*-test, and Cohen's *d* for mean difference. The study deemed to be significant because of the possible relatedness of the gametophyte development to its nitrogen-absorbing ability which

confers the capability to remedy polluted water. Freehand cross sections of the anther, ovary and seed were prepared and stained with acetocarmine, safranin, and iodine. Banderang español androecium consists of two trimerous whorls: the epipetalous staminate inner whorl and the episepalous outer whorl. Septa homogenization and stomium opening characterize the maturation of the bisporangiate anther. In microsporogenesis, the microsporocysts undergo successive cytokinesis, callose deterioration, and mitosis to form a vacuolated microspore giving rise to vegetative nucleus and two sperm cells. Pollen germinates best at 20% sucrose peaking on the 24th hour (*d* = 0.91). *C. x generalis* ovary follows the monosporic polygonum development. A megasporocyte divides meiotically giving rise to a megaspore tetrad with the only functional megaspore undergoing mitosis thrice to form an octanucleated gametophyte. Morphometric analysis revealed a positive correlation between the anther size and the ovary size. While petal size and sepal size are proportional, they pose no significant relationship to the staminode growth. Also, anther growth peaks at the length of 1.71 cm (± 0.4 cm). Recommendations for further studies include usage of other histochemical stains and through study of embryogeny.

Prof. Emmanuel D. Delocado, Department of Biology, School of Science and Engineering, Ateneo de Manila University, Loyola Heights, Quezon City 1108, Philippines

Using Ecological Studies about Naturalization Rate of Alien Plants in Teaching Biology

Ji-Hyon Kil¹, Kew-Cheol Shim²

¹*National Institute of Environmental Research,*

²*Kongju National University, South Korea*

Invasive alien species are one of main factors to affect on indigenous plants, ecosystems and biodiversity. After surveying plants, which habituated in some areas, both indigenous and alien plants were listed and naturalization rate of alien plants (NRAP) were calculated. Study sites are railway and riparian ecosystems, where could be often mediated to by transport, animals and humans. Collecting data were analyzed comparatively to examine ecological implications. The use of NRAP can be an effective tool to understand natural

environment ecologically and ecological impacts of invasive alien plants. Challengeable topics and well-structured experiments for students could derive students actively to engage and to motivate their curiosities. Ecological studies should induce scientific meaning related to topics, experimental experiences and then encourage them to invent original ideas. Practical works based on ecological studies have potentials to facilitate scientific inquiry activities.

Prof. Kew-Cheol Shim, Department of Biology Education, Kongju National University, Gongju, Chungnam, South Korea

Scanning Electron Microscope (SEM)-aided Screening of Wild Type *Bacillus thuringiensis* Parasporal Crystalline Protein-based on Structure against *Aedes aegypti* Larvae

**Jing R. Bautista, Faith B. Amorado,
Maria Luria S. Orbita, Franco G. Teves**
MSU-Iligan Institute of Technology, Philippines

Dengue has claimed thousands of Filipino lives in the past few years with incidence that defy the traditional seasonality of this mosquito-borne viral infection. Preventive measures include destruction of known mosquito habitats, use of mosquito nets and house screens, fogging and use of chemical insecticides. While adults may be killed or prevented from spreading the virus through bites, the eggs and larvae can survive and continue to spread the virus to the human population. To date, there are only very few identified strains of the bacterial species *Bacillus thuringiensis* that produce the parasporal crystalline protein effective against Dipteran insects such as mosquitoes. Moreover, there seems to be a relationship between protein crystal shape and its efficiency as larvicidal protein. The gold standard for bacterial identification is considered to be 16S rRNA sequencing. However, in this study, SEM was employed to determine the parasporal crystalline protein structure and deduce its larvicidal strength through actual *Aedes aegypti* larval toxicity assay. The method is fast, straightforward, and comparably cheaper than the alternative molecular technique in identifying candidate bacterial strains as sources of bioinsecticides against the dengue-carrying mosquito *A. aegypti*. It was also demonstrated in this study that a single bacterial strain could produce differ-

ent crystal shapes corresponding to characteristic LC50 (lethal concentration that kills 50% of the mosquito larvae). This is the first SEM study on *B. thuringiensis* parasporal crystalline protein in the Philippines as far as literature is concerned.

Prof. Jing R. Bautista, Department of Biological Sciences, College of Science and Mathematics, MSU-Iligan Institute of Technology, Iligan City, Philippines

Ethnobotany of Ferns and Fern Allies in Mount Macabol Marilog District Davao, Philippines

Adorico M. Aya-ay
University of the Immaculate Conception, Philippines

Ethnobotany of ferns and fern allies was conducted in an indigenous community at Mt. Macabol, Barangay Salaysay, Marilog District, Davao City, to document the tribal uses of the different species of pteridophytes in the forest. The gathering of data was done through interview with the chosen members of the community and through the collection of herbarium specimens. Seventy-one species of ferns and fern allies are collected. Thirty-seven species (52%) have uses in the tribe. These useful species serve wither as food, medicine, ornament, ceremonial materials, or in other forms of uses for the community. This paper presents the local/common names, scientific names and the specific uses of the plants. The data obtained show that despite human encroachment, Mt. Macabol remained a home to a good number of ferns and fern allies useful to the residents.

Prof. Adorico M. Aya-ay, University of the Immaculate Conception, Bonifacio Street, Davao City, Philippines

The Distribution and Relative Abundance of Medium-sized Arboreal Mammals at Mount Mahuson, Mount Apo Key Biodiversity Area

Adorico M. Aya-ay
University of the Immaculate Conception, Philippines

Field surveys of arboreal mammals were conducted in the forests of Mt. mahuson in Arakan Valley, North Cotabato Province from October 2009 to March 2011. With concentration on the prey items of the Philippine eagle *Pithecophaga jefferyi*, the team searched for

Mindanao flying lemur *Cynocephalus volans*, Mindanao flying squirrel *Petinomys crinitus*, palm civet *Paradoxurus hermaphroditus*, and long-tailed macaque *Macaca fascicularis*. The team conducted mammal surveys for a total of 48.38 observation hours in seven contiguous line transects (with a total length of 14 km). The team was also able to gather data from opportunistic sampling and dialogues with parabiologists. Results indicate that among these species, the long-tailed macaques are most abundant particularly in the remaining forests at lower altitude (below 1300 masl), followed by the Mindanao flying lemur and palm civet. The Philippine tree squirrel scored only “frequent” whereas both the Mindanao flying squirrel and pygmy squirrel were found rare in the study area. The greater number of individuals encountered in those particular forests at lower altitude could be attributed to the greater quality of (i.e. thicker canopy cover, larger trees) habitats in the area. The greater abundance of macaques relative to lemurs in the area suggests that the lemurs are not the primary prey items of the nesting pair in Mt. Mahuson. This is contrary to the fact that a large bulk of prey items of the Philippine eagle population in Mindanao consists of flying lemurs. Future research should focus on testing this hypothesis in order to further assess the prey population and foraging behavior of the eagle pair in the mountain.

Prof. Adorico M. Aya-ay, University of the Immaculate Conception, Bonifacio Street, Davao City, Philippines

Vegetation Analysis of the Pasonanca Natural Park in Zamboanga City

Jocelyn D. Partosa, Jamaica Lota
Ateneo de Zamboanga University, Philippines

This study analyzed the vegetation of the Pasonanca natural Park in Zamboanga City. The sampling sites covered Canucutan, Muruk 3 and Baluno. Three stations were established at intervals of 300 m each in Canucutan and Muruk 3 and four stations in Baluno. Five nested quadrats, each measuring 100 m² with subplots measuring 5 m × 5 m and 1 m × 1 m were plotted at intervals of 10 m in each station. There were 283 individuals belonging to 45 species, 27 genera and 25 families. The taxonomic composition is mainly angiosperms and pteridophytes comprising 82.2% and

17.8%, respectively. There are 30 species with known geographic range; and of these, 60% are endemic or native, 33.3% are cosmopolitan and 6.7% are exotic. Data on species composition, distribution, importance values, and diversity indices all suggest that species diversity is high in all three sites. Species evenness though was fairly high in Canucutan compared to Muruk 3 and Buluno based on Simpson's measure of evenness. Conversely, all three sites have comparatively high species evenness based on Shannon-Wiener's index. There are few species under the threatened category (either vulnerable or critically endangered). Several species though are either not threatened (33%) or not evaluated (27%). The ecologically important species belong to families Araceae (*E. pinnatum* and *Rhapidophorra* sp.), Arecaceae or Palmae (*Arenga* sp., *Calamus* sp. And *Caryota* sp.), Euphorbiaceae (*M. tanarius*), Maranthaceae (*D. canei-formis*) and Athyriaceae.

Prof. Jocelyn D. Partosa, Ateneo de Zamboanga University, Zamboanga City, Philippines

Assessment of the Mangrove Forest Structure of Selected Marine Protected Areas of Southeast Cebu

Cristina C. Genosa, Delia P. Buctuan, Rhodora Z. Layumas, Ida L. Ssantiago, Patricia M. Mailed
Southwestern University, Philippines

A study was conducted to assess the mangrove forest structure of selected Marine Protected Areas of Southeast Cebu using the descriptive survey method. Specifically, the study sought to identify the mangrove species composition, density and describe the community types and zonation of the selected study sites. There were five identified study sites which included: (1) Brgy. Bagacay Marine Fish Sanctuary, Sibonga; (2) Sitio Pungtod, Siala, Sibonga; (3) Brgy. Taloot, Argao; (4) Brgy. Cawayan Marine Sanctuary, Dalaguete; and (5) Sitio Nonoc, Brgy. Daan-lungsod, Alcoy. The study identified 19 mangrove species which included 12 major mangrove species, four minor mangrove species and three mangrove associate species. The densest tree stand was located in site (3) followed by site (1), (4), (2) and the least stand density as in site (5). Summing all densities from all sites for each species, it

was found out that the most dominant major mangrove species was *Rhizophora* sp., followed by *Avicennia* sp. and *Sonneratia* sp. These three species are characterized as highly adapted to sandy/muddy substrates which is the most common feature of all the study sites. All study sites were categorized as riverine forest types which are tall flood plain forests along flowing waters such as tidal rivers and creeks. Conditions in this type of forest are favorable for extensive growth due to flushing by daily tides.

<Sorry for missing some of the last sentences.>

Prof. Cristina C. Genosa, Southwestern University, Cebu City, Philippines

Seed Morphology of Two Endemic Species of *Medinilla* Gaud (Melastomaceae): Its Implications for Disposal and Germination

Jan Lorie M. Robil, Vivian S. Tolentino
Ateneo de Manila University, Philippines

The morpho-anatomy of seeds of *Medinilla magnifica* and *Medinilla* sp. was studied under light and scanning electron microscope (SEM). The goal of this study is to delineate the morpho-anatomical characteristics of seeds of two closely resembling *Medinilla* spp. Aside from acrodromous leaf venation, melastomaceae is recognized among dicots with their numerous exalbuminous seeds. This is true among *Medinilla* species, where seeds develop from anatropous ovules of the epigynous ovary of the flower. Based on the close resemblance of the two plants in terms of vegetative and reproductive anatomy, one would expect that the seeds of the plants would also be similar in terms of their form. However, the investigation revealed that there are several notable differences in their form, especially in their external features. In terms of shape, mature seeds of *M. magnifica* are round oval as compared to those of *Medinilla* sp. Which are rather flattened toward the micropylar end. A strikingly remarkable difference was observed between the surface textures of the seeds under SEM. The seeds of *Medinilla* sp. Has rather smooth surface due to tabular cells forming jigsaw puzzle-like pattern. The seed surface of *M. magnifica*, on the other hand, has a relatively rougher surface due to cells forming blister-like projections. The seed coat surface has been studied to

have significant role in seed dispersal by biotic and abiotic factors. The rough seed surface of *M. magnifica* could be of great advantage when it comes to adherence to aerial substrates during dispersal while the smooth seed surface of *Medinilla* sp. Could be adaptive to dispersal by animal frugivores. With regards to internal anatomy, particularly to embryo form and position, the seeds of two plants are relatively similar conforming to the general characteristics of Melastomaceae. The short embryos of the seeds are straight or slightly bent, lying along the long axis. At longitudinal section, the lignified seed coat is thicker in the chalazal end of the seed and diminishing towards the micropylar end. The form and position of the embryos are reflective of the phanerocotylar germination of the seeds, which is common in the family. In conclusion, the seeds of *Medinilla* sp. and *M. magnifica* have distinctly different external features but have relatively similar internal anatomy. The observed morpho-anatomical differences and similarities of the seeds reflect morpho-functional and developmental adaptations, which are important for dispersal and establishment of the plants. These observations are of great consideration for conservation of the plants since *M. magnifica* and *Medinilla* sp. are considered endangered and rare, respectively.

Prof. Jan Lorie M. Robil, Department of Biology, School of Science and Engineering, Ateneo de Manila University, Loyola Heights, Quezon City 1108, Philippines

GIS Mapping on Invasive Aquatic Plant *Eichhornia crassipes* (Commelinales: Pontedericeae) among Wetlands and Farmlands in Carcar, Cebu, Central Philippines

Alejandro C. Manlangit, Jr.
University of San Carlos and Carcar Central NHS – Carcar City Division, Philippines

Aquatic invasive plant species are widely considered to be one of the greatest threats to biodiversity. Water hyacinth (*Eichhornia crassipes*) is the subject of the study on its invasive nature. The objective is to create a GIS database to model for management of its occurrence, verification, and control. The data included both physical and environmental parameters surveyed

in the field specifically its location and coordinates. There are ten villages surveyed having wetlands and its corresponding coordinates through GPS counter. Research study points were identified prior to actual recordings. Quadrat methods were established randomly to determine species relative density. Elevation varies from 0.7 – 88 m above sea level. Station 6 has the highest percentage covered having 41.41% and Station 1 having 3.38% as the lowest. Relative density and evenness was achieved highly at Station 2 - 5 having 281 species per 10m² and Station 2 - 4 has the lowest record having 102 per 10m²

Prof. Alejandro C. Manlangit, Jr., Biology Department, University of San Carlos, Cebu City, and Carcar Central NHS – Carcar City Division, Cebu, Philippines

Comparative Developmental Morpho-anatomy of Idioblastic Oil cavity in Stem, Leaf and Rind of *Citrus x sinensis* ‘Satsuma’ (L.) Osbeck (Sweet Orange) and *C. x microcarpa* Bunge (Calamansi)

Emmanuel D. Delocado, Josh Nathan L. Ngai, Carol Angeline G. Yu, Reshan D. Baoas, Sol Kim, Vivian S. Tolentino

Ateneo de Manila University, Philippines

Five years ago, the international market with crisis as prices of petroleum products fuelled up by almost 600%. Because of this, the search for an alternative energy source among local species has been a fad. This study investigates the development of idioblastic oil cavities in the stem, leaves, and fruit rind of *Citrus x sinensis* (L.) Osbeck (sweet orange) and *C. x microcarpa* Bunge (calamansi). Data from this study serves as a progeny study for oil extraction strategies and as a contribution to the growing literature on oil cavity development. Freehand cross sections of the stem, leaf and fruit pericarp for both plants were prepared and stained using Nile Blue sulphate, Sudan IV, and Sudan Red. Oil cavities originate as parenchymal oil cells which differentiate into polyhedral inner cells and flattened boundary cells connected to the epidermis with a copule. Eventually, the flattening and swelling of inner cells leads to lysigenic development of oil cavities. Oil particulates start moving to the cavity periphery and attract possible remaining oil substances

in the now flattened inner cells. The ontogeny of the idioblastic oil cavity culminates with the oil deposit forming in the cavity. Three layers of cell wall are observed on rind oil cavity of both plants: a suberin layer sandwiched between two cellulose layers. Oil deposits stain positively in all three histochemical stains. ANOVA (95% confidence level) reveals a positive correlation between the diameter of the *C. x sinensis* fruit and the diameter of the oil cavities. In addition, the diameter of oil cavity of *C. x microcarpa* increases significantly as it matures, then stop after it ripens. Recommendations for further studies include the preparation of fresh and permanent sections, morpho-anatomical investigations of root and flowers, the usage of other histochemical stains, and an exploration on oil extraction techniques.

Prof. Emmanuel D. Delocado, Department of Biology, School of Science and Engineering, Ateneo de Manila University, Loyola Heights, Quezon City 1108, Philippines

Two New Generic Records in Philippine Rubiaceae Based on Plastid and Nuclear DNA Sequence Data

Axwl H. Arriola^{1,2}, Grecebio Jonathan D. Alejandro²

¹University of the East, ²University of Santo Tomas, Philippines

The tribe Vanguerieae (Rubiaceae) in the Philippines is represented by *Canthium* (20 spp.) and *Psydrax* (monotypic). Recent studies done in the tribe showed that most of its representatives form a heterogenous assemblage. The genus *Canthium* was restricted by having supraaxillary spines. This raised questions on the position of the PPhilippine *Canthium* as members are without spines. In this first molecular study of two Philippine *Canthium* (*C. monstrosum* and *C. ramosii*), *trnL-F* and ITS regions were sequenced, assembled and aligned manually using Se-Align v2.0 and subsequently analyzed using MrBayes 3.1.2. Interestingly, the majority-rule consensus tree revealed that the two Philippine species were nested in two different clades with high support; *C. monstrosum* within the *Keetia* clade (PP = 0.92) while *C. ramosii* grouped with the *Pyrostria* clade (PP = 0.98). Therefore, we proposed two new combinations, the *Keetia monstrosum* (A. Rich.) Arriola & Alejandro and *Pyrostria ramosii*

(Merr.) Arriola & Alejandro. This study establishes for the first time the two genera (*Keetia* and *Pyrostria*) in the Philippines. The IUCN red list declared that conservation status of *Keetia* and *Pyrostria* were from vulnerable to critically endangered.

Dr. Grecebio Jonathan D. Alejandro, College of Science and Research Center for Natural and Applied Sciences, University of Santo Tomas, Espana, Manila 1015, Philippines

Reproductive Characteristics of *Grapsus tenuicrustatus* (Herbst, 1783) (Decapoda: Grapsidae) Found in Talim Bay, Lian, Batangas, Philippines

Michael A. Clores¹, Gliceria Ramos²

¹Ateneo de Naga University, ²De La Salle University, Philippines

Understanding the breeding potential of a well dispersed marine invertebrates, like crabs, entails determining variation of its productive characteristics (Duran, 1991). The present study determined some reproductive characteristics of *Grapsus tenuicrustatus* (Herbst, 1783), a brachyurans (true crabs) which are among the most widespread and diverse groups of invertebrates. Samples of *G. tenuicrustatus* were collected at Talim Point (130° 57' 55.43", 1200 36' 20.36"E), a portion of Talim Bay at Barangay Ligtasin, Lian, Batangas, at rocky areas near the seagrass beds. Results revealed that there were more male *G. tenuicrustatus* (n = 81) than female (n = 52) with a sex ratio of 1.13:1. Thirty percent of the samples were ovigerous females. Ovigerous female crabs have the largest carapace length (CL) (mean ± SE: 31.25 ± 1.43) compared with the males (mean ± SE: 22.14 ± 0.73) and non-ovigerous females (mean ± SE: 26.63 ± 1.12). Based on one-way ANOVA, the CL differences were significant (F = 20.383, df = 2, p < 0.01). Difference in CL between non-ovigerous and ovigerous females was also found to be significant based on t-test for independent samples (t = 2.582, df = 70, p < 0.05). There was a non-conspicuous bi-modal size distribution for all the crabs, with non-normal distributions for all crabs (one-sample Kolmogorov-Smirnov (KS) = 1.735, p < 0.05) and for males (KS = 1.464, p < 0.05), but not when all females or ovigerous females only were grouped together. The size-frequency distribu-

tions of males and females are significantly different from each other (KS = 2.582, p < 0.0001). The same was observed between the size frequency of ovigerous and non-ovigerous female crabs (KS = 1.502, p < 0.05). Moreover, there were more ovigerous crabs belonging to the first year age class (CL = 16 – 33) (53%) than those that belong to the older class (CL = 34 – 43) (47%). The fecundity of *G. tenuicrustatus* ranged from 4400 (CL = 16 mm) to 26,400 (CL = 43 mm) eggs. Egg volume ranged from 0.40 ml to 2.40 ml. <Sorry for missing some of the last sentences.>

Michael A. Clores, Department of Natural Sciences, Ateneo de Naga University, Philippines

Anti-bacterial Potential of the Cultivated and Wild Type Adlai (*Coix lacryma-jobi* Linn.) Plant Extracts

Jeany Cuaresma Dupo, Lorenza G. Lirio

Benguet State University, Philippines

Antibacterial potential of aqueous, ethanolic and methanolic extracts of the different plant parts of the cultivated and wild types of "adlai" (*Coix lacryma-jobi* Linn.) was investigated against *Staphylococcus aureus* and *Escherichia coli*, gram-positive and gram-negative bacteria, respectively. The antibacterial activity was performed by paper disc-diffusion method. The presence of antibacterial agents in the plant extract was determined by phytochemical analysis and thin layer chromatography. The zones of bacterial inhibition produced by the test organisms were subjected to analysis of variance and the differences in treatment means by DMRT at P=0.05. No activity was seen in the aqueous extracts while a slight antibacterial activity was observed on the ethanolic and methanolic stem and root extracts of both plant types except the methanolic stem extract (50mg/ml) of the cultivated type which showed a very significant growth of inhibition against *S. aureus* (14.5mm) and *E. coli* (13.2mm). The antibacterial activity was comparable with the standard antibiotic, streptomycin (15.2mm). Results of the phytochemical studies revealed the presence of alkaloids, steroids, flavonoids, saponins and polyphenols which were active against both gram-positive and gram negative-bacteria. Moreover, thin-layer chromatography confirmed the presence of sterols, phenols and essential

oils which is rich in triterpenes in the methanolic extract of the stem (cultivated type). The ethyl acetate:methanol (95:5) showed the greatest resolution with 4 spots using different detection method and is recommended as the mobile phase which provides a R_F value of 0.59 – 0.76. The results obtained in the present study suggest that the plant can be used in treating diseases caused by the test organisms and provide very important information to the medical and pharmaceutical industries in developing cheaper drugs of plant origin.

Prof. Jeany Cuaresma Dupo, Benguet State University. La Trinidad, Benguet, Philippines

A Study on the Anti-Diabetic Potential of Pansit-Pansitan (*Peperomia pellucida*) Extract

**Patricia Anne P. Bonifacio¹, Jason C. Alcares¹,
Edna A. Amparado²**

¹*Philippine Science High School – Main Campus,*

²*University of the Philippines, Diliman, Philippines*

The use of pansit-pansitan (*Peperomia pellucida*) extract as a substitute treatment to high blood glucose levels of diabetic patients was verified to provide a more affordable alternative to diabetic maintenance medications. One kilogram of leaves was acquired and air-dried for two weeks, then steeped in 99% ethanol for three nights. The pure extract was obtained through a rotary evaporator then air-dried for one day. Alloxan was initially supplemented at a 150 mg/kg-body weight dosage to four groups of mice with three replicates per setup to induce diabetes. Controlled variables include environmental conditions and diet, while experimental variables consist of extract concentration and blood glucose levels. The 50.00 mg/kg-body weight extract was prepared in ethanol at 50 – 75% concentrations then administered to two setups on non-diabetic mice. Glucose levels were checked after the first hour, then after every four hours for 24 hours. Preliminary data showed significant lowering in the levels of mice treated with extract solution. The ANOVA tests confirmed this initial conclusion. The success of this study presents a cheaper alternative diabetic treatment to underprivileged patients. A new source of income may also be ascertained if the me-

dicinal potential of this common plant is further investigated.

Patricia Anne P. Bonifacio, Philippine Science High School – Main Campus, Quezon City, Philippines

<Workshops>

Create a Sheep Mascot While Feeling the Texture of Wool

Hirofumi Naekawa

Tokyo Gakugei University, Japan

Despite sheep is a familiar animal for people, we believe that the number of children and students who have never seen or touched the animal in their life has been increasing in recent years. This workshop is an experience-based booth where you can actually touch wool and create a sheep mascot while feeling the texture of wool. I will exhibit how to make the mascot. You can bring the sheep mascot back to home. As the mascot materials for one person, you only need one gram of wool, two laces with wire 24 cm in length, and scissors.

Dr. Hirofumi Naekawa, Department of Biology, Tokyo Gakugei University, Tokyo, Japan

Integrating Environmental Education into Integrated Science Courses in the K-12 Curriculum – A Workshop on Adapting Ideas from Native Seasons, Project Wet, Wild and Learning Tree into the Filipino Context

John Paolo Dalupang

Ateneo de Manila University, Philippines

With the implementation of the new K-12 curriculum in the Philippines, educators in various levels in Basic Education have raised concerns about challenges in implementing an all-integrated and spiraling approach in teaching science. This concern is most resonant in the 7-10 grade levels where concepts in the major science courses, namely biology, chemistry, and physics, are taught in the same grade levels within the same school year. While textbooks and instructional materials are still being developed for this new curriculum, teachers are left with their creativity and ingenuity on how to effectively cover these topics.

This workshop will provide educators in various levels with ideas that can help with creating learning resources and lessons that can help with the issue of integration. The main ideas that will be presented in the workshop will come from established environmental education curricula that have been used in the United States for over 10 years – Native Seasons, Project WET, WILD, and Learning Tree. Through developed in another country, the activities presented in these learning materials are very flexible and can be easily adapted to the context of its learners. Grounded in environmental education, these curricula used what's available in the learner's community environment to understand the concepts in science. The workshop will begin with a brief description of the tenets of environmental education and the background of each of the curricula. The rest of the session will consist of hands-on and lively activities that are mostly geared to some topics in the new Philippine K-12 Grade 6, 7, and possibly Grade 8 (if time permits) science curricula in science. Attendees from other countries might want to prepare a list of topics for their Grade 6- or 7-equivalent levels for referencing with the learning materials. The proponent of this workshop neither in any way represent the organizations that developed the mentioned learning materials nor is connected with the same organizations. The proponent just wants to share options that became useful during his teaching career. It is a workshop where teachers can share

ideas to help cope with the demand of the new curriculum.

Prof. John Paolo Dalupang, Ateneo de Manila University, Quezon City, Philippines

A Simple Laboratory Work on DNA Extraction at Secondary School Level

Yoshihiko Yonezawa, Noriko Banba
Naruto University of Education, Japan

For a biology class at the secondary school level, a simple method for DNA extraction from flower buds of broccoli was developed using a mixture of 4% NaCl solution and 1% SDS (sodium dodecyl sulfate) solution (24:1) as an extraction medium. However, this simple method is unsuitable for the DNA extraction from fruits of banana and strawberry, because the UV absorption spectrum of the ethanol precipitate from the extracts of banana and strawberry did not show the typical absorption curve of DNA. Therefore, we propose broccoli, which can be obtained at any time and by the reasonable cost, as a plant material. In addition, the diphenylamine-colorimetric method is also proposed for detecting extracted DNA. In this workshop, we demonstrate this simple method for DNA extraction from flower buds of broccoli.

Prof. Yoshihiko Yonezawa, Department of Biology, Naruto University of Education, Tokushima, Japan

Publication

A book, *Biology Education for Social and Sustainable Development*, was published in 2012 by Sense Publishers, Rotterdam, Netherlands. Some papers presented at the **23rd Biennial Conference of the AABE** which was held in Singapore in October 2010 were compiled in this book by the co-organizers of the conference, Dr. Mijung Kim and Dr. C. H. Diong. You can refer to the abstracts of these papers in **the sixth volume of the Asian Journal of Biology Education** (2012).
